arXiv:1102.1872v4 [math.NT] 16 Nov 2011

ON SOME ARITHMETIC PROPERTIES OF AUTOMORPHIC FORMS OF GL_m OVER A DIVISION ALGEBRA

H. GROBNER AND A. RAGHURAM

Abstract. In this paper we investigate arithmetic properties of automorphic forms on the group $G' = GL_m/D$, for a central division-algebra D over an arbitrary number field F. The results of this article are generalizations of results in the split case, i.e., $D = F$, by Shimura, Harder, Waldspurger and Clozel for square-integrable automorphic forms and also by Franke and Franke-Schwermer for general automorphic representations. We also compare our theorems on automorphic forms of the group G' to statements on automorphic forms of its split form using the global Jacquet-Langlands correspondence developed by Badulescu and Badulescu-Renard. Beside that we prove that the local version of the Jacquet-Langlands transfer at an archimedean place preserves the property of being cohomological.

Contents

1. Introduction and statements of results 1
2. The general linear group and its inner forms 4
3. Generalities on automorphic representations of $G'(\mathbb{A})$ 6
4. A classification of the cohomological, irreducible, unitary dual of $GL_k(\mathbb{H})$ and $GL_n(\mathbb{R})$ 8
5. The local and global Jacquet-Langlands transfer and its interplay with cohomology 13
6. Algebraic and regular algebraic representations of $G'(\mathbb{A})$ 18
7. Spaces of automorphic cohomology and rational structures 22
8. Results on rationality fields 29
References 33

1. Introduction and statements of results

Let D be a central division algebra of index d over a number field F. The group $G' = GL_m/D$ of invertible $m \times m$ - matrices with entries in D defines a connected, reductive group over F and is an inner form of the split general linear group $G = GL_n/F$, $n = dm$. In the split case, i.e., if $D = F$, many important results on the arithmetic theory of automorphic forms are known due to several people. Within the scope of the present paper, for cusp forms one should particularly mention the work of Shimura [35] for $n = 2$ and F totally real, Harder [15] and Waldspurger [41] for $n = 2$ and any F, Clozel [8] for general n and F; and for general automorphic forms of the group G we mention Franke [11] and Franke–Schwermer [12]. In [41] one may also find results in the very special non-split case $m = 1$ and $d = 2$.

The main aim of this article is to study the arithmetic of automorphic forms on G' and to generalize some of the results of the above mentioned people to the case of GL_m/D for a general m and a general D.

Date: January 12, 2013.

2010 *Mathematics Subject Classification.* Primary: 11F70, 11F75, 22E47; Secondary: 11F67.

Key words and phrases. Inner forms of GL_n, global Jacquet-Langlands, algebraic representations, regular algebraic representations, cuspidal automorphic forms, rationality properties, $A_{\mathfrak{q}}(\lambda)$-modules, unitary dual.

H.G. is supported by the Austrian Science Fund (FWF) Erwin Schrödinger grant, J 3076-N13. A.R. is partially supported by the National Science Foundation (NSF), award number DMS-0856113, and an Alexander von Humboldt Research Fellowship.

For a central division algebra D over F and $m \geq 1$, Badulescu [1] and Badulescu–Renard [2] have recently proved the existence of the global Jacquet-Langlands transfer JL from discrete series automorphic representations of $G'(\mathbb{A})$ to discrete series automorphic representations of $G(\mathbb{A})$. This establishes a special instance of Langlands functoriality and forms an important instrument for the analysis of arithmetic properties of square-integrable automorphic forms of $G'(\mathbb{A})$. We also explicitly describe the interplay of the local Jacquet-Langlands transfer and cohomology.

Our first theorem, which is of an arithmetic nature, deals with the notion of a regular algebraic representation of $G'(\mathbb{A})$. In the split case, i.e., $D = F$, Clozel gave a definition of a representation being regular algebraic: A discrete series automorphic representation Π of $G(\mathbb{A})$ is called algebraic, if each of the irreducible representations in the archimedean part Π_∞ corresponds via the local Langlands parametrization to a sum of n algebraic characters of $\mathbb{C}^*$. An algebraic representation Π is furthermore called regular, if the infinitesimal character of Π_∞ is regular. It easily follows from Clozel [8] that a cuspidal automorphic representation Π is regular algebraic if and only if Π_∞ is essentially tempered and cohomological with respect to a certain coefficient system E_μ.

In this paper we extend Clozel's notion as follows: We call a discrete series automorphic representation $\Pi' = \Pi'_\infty \otimes \Pi'_f$ of $G'(\mathbb{A})$ algebraic (resp., regular algebraic) if its global Jacquet–Langlands transfer $\Pi = JL(\Pi')$ is. With this definition we prove the following generalization of Clozel's result, cf. Thm. 6.9.

Theorem 1.1. *Let Π' be a discrete series automorphic representation of $G'(\mathbb{A})$ and assume that $JL(\Pi')$ is cuspidal. Then the following are equivalent:*

(i) *Π' is regular algebraic.*

(ii) *Π'_∞ is cohomological and essentially tempered.*

Furthermore, the implication (ii)⇒(i) does not seem to need the assumption that $JL(\Pi')$ is cuspidal. See Remark 6.10. In contrast, the implication (i)⇒(ii) may fail without assuming $JL(\Pi')$ is cuspidal, cf. Ex. 6.11.

Given our definition of an algebraic representation, we also generalize Clozel's "Lemme de pureté" to the case of G', cf. Lem. 6.12:

Lemma 1.2 (Purity Lemma for G')**.** *If Π' is an algebraic representation of $G'(\mathbb{A})$ and $JL(\Pi')$ cuspidal, then there is a $\mathsf{w} \in \mathbb{Z}$ such that for all archimedean places v, the algebraic characters of $\mathbb{C}^*$ associated to $JL(\Pi')_v| \cdot |_v^{\frac{1-n}{2}}$ are of the form $z \mapsto z^p(\overline{z})^q$ with $p + q = \mathsf{w}$.*

Again, we give an example that the cuspidality assumption on $JL(\Pi')$ cannot be removed from the statement of the lemma, see Ex. 6.13.

Next, we analyze the interplay of cohomological automorphic forms of $G'(\mathbb{A})$ and the action of $\mathrm{Aut}(\mathbb{C})$. Following Waldspurger [41] and Clozel [8], for any group representation ν and $\sigma \in \mathrm{Aut}(\mathbb{C})$, there is the σ-twisted representation ${}^\sigma\nu := \nu \otimes \mathbb{C}_{\sigma^{-1}}$. In particular, this definition applies to the finite part Π'_f of an automorphic representation Π of $G'(\mathbb{A})$. The action of $\sigma \in \mathrm{Aut}(\mathbb{C})$ on the highest weight module E_μ is via its permutation action on the embeddings of F into $\mathbb{C}$. (See, for example, Raghuram–Shahidi [29, Sections 3.1 and 3.3].) Now, let

$$S_{G'} = G'(F)\backslash G'(\mathbb{A})/K'^{\circ}_\infty,$$

where K'°_∞ is the topological connected component of the product of the center of G'_∞ and a maximal compact subgroup of G'_∞. A finite-dimensional highest weight module E_μ defines a sheaf $\mathcal{E}_\mu$ on $S_{G'}$. The corresponding sheaf cohomology $H^q(S_{G'}, \mathcal{E}_\mu)$ can be computed using Betti-cohomology and hence for any $\sigma \in \mathrm{Aut}(\mathbb{C})$, there is a σ-linear isomorphism

$$\sigma^* : H^q(S_{G'}, \mathcal{E}_\mu) \xrightarrow{\sim} H^q(S_{G'}, {}^\sigma\mathcal{E}_\mu).$$

On the other hand, $H^q(S_{G'}, \mathcal{E}_\mu)$ is isomorphic to the $(\mathfrak{g}'_\infty, K'^{\circ}_\infty)$-cohomology of the space of automorphic forms on $G'(\mathbb{A})$, which follows from Franke [11], and inherits from that a decomposition of $G'(\mathbb{A}_f)$-modules

$$H^q(S_{G'}, \mathcal{E}_\mu) \cong \bigoplus_{\{P'\}} \bigoplus_{\varphi_{P'}} H^q_{\{P'\},\varphi_{P'}}(G', E_\mu),$$

cf. [12] and [27]. Here, the sums are ranging over the associate classes of parabolic F-subgroups $\{P'\}$ of G' and attached associate classes of cuspidal automorphic representations $\varphi_{P'}$ of the corresponding Levi subgroup L'. In particular, the summand indexed by $\{G'\}$ gives the cohomology of the space of cuspidal automorphic forms of $G'(\mathbb{A})$, which is usually called the cuspidal cohomology of G'.

We prove the following result, cf. Thm. 7.22, which says that for regular highest weights, σ^* respects this fine decomposition into cuspidal supports.

Theorem 1.3. *Let E_μ be a regular highest weight representation of G'_∞ and $\sigma \in \mathrm{Aut}(\mathbb{C})$. For each associate class of parabolic F-subgroups $\{P'\}$, and each associate class of cuspidal automorphic representations $\varphi_{P'}$, the summand $H^q_{\{P'\},\varphi_{P'}}(G', E_\mu)$ of the global cohomology group $H^q(S_{G'}, \mathcal{E}_\mu)$, is mapped by σ^* isomorphically onto the summand $H^q_{\{P'\},{}^\sigma\varphi_{P'}}(G', {}^\sigma E_\mu)$ of $H^q(S_{G'}, {}^\sigma\mathcal{E}_\mu)$ for a unique associate class ${}^\sigma\varphi_{P'}$:*

$$H^q_{\{P'\},\varphi_{P'}}(G', E_\mu) \xrightarrow[\sigma*]{\cong} H^q_{\{P'\},{}^\sigma\varphi_{P'}}(G', {}^\sigma E_\mu).$$

If $H^q_{\{P'\},\varphi_{P'}}(G', E_\mu) \neq 0$ and a representative P' in the associate class $\{P'\}$ has Levi factor L' and $\varphi_{P'}$ is represented by a cuspidal automorphic representation Π' of $L'(\mathbb{A})$, then $\Pi' \otimes \rho_{P'}$ is cohomological. The σ-twist ${}^\sigma\Pi'_f \otimes {}^\sigma\rho_{P'_f}$ of its finite part is the finite part of a unique cohomological cuspidal automorphic representation Ξ' and the associate class ${}^\sigma\varphi_{P'}$ is uniquely determined by the representation $\Xi' \otimes \rho^{-1}_{P'}$.

Under the assumption of the regularity of E_μ, this theorem is a generalization of the analogous result of Franke–Schwermer [12] in the split case, $D = F$. In this setup, it is also a simultaneous generalization of Franke's [11, Theorem 20] on the compatibility of σ^* with the $\{P\}$-decomposition and Clozel's [8, Théorèm 3.13], which states that the σ-twist ${}^\sigma\Pi_f$ of the finite part of a cohomological cuspidal automorphic representations Π of $G(\mathbb{A})$ is the finite part of a cohomological, cuspidal automorphic representation.

If we consider the action of $\mathrm{Aut}(\mathbb{C})$ on a regular algebraic representation and ask whether it is compatible with the global Jacquet-Langlands transfer JL, then we obtain the following result, cf. Thm. 7.27:

Theorem 1.4. *Let Π' be a regular algebraic representation of $G'(\mathbb{A})$ and assume that $JL(\Pi')$ is cuspidal. Then for all $\sigma \in \mathrm{Aut}(\mathbb{C})$, ${}^\sigma\Pi'_f$ is the finite part of a representation Ξ' which is regular algebraic and $JL(\Xi')$ is cuspidal.*

In the final section we also prove some arithmetic results on the rationality field

$$\mathbb{Q}(\Pi'_f) = \{z \in \mathbb{C} \mid \sigma(z) = z \quad \forall \sigma \in \mathrm{Aut}(\mathbb{C}) \text{ for which } {}^\sigma\Pi'_f \cong \Pi'_f\}$$

of the finite part of a cuspidal automorphic representation Π' of $G'(\mathbb{A})$ which is regular algebraic and lifts to a cuspidal automorphic representation $JL(\Pi')$. At infinity, define $\mathbb{Q}(\mu)$ as the fixed-field in $\mathbb{C}$ of the group of automorphisms of $\mathbb{C}$ which fix a highest weight representation E_μ with respect to which Π'_∞ has non-zero $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-cohomology. Now, let the field $\mathbb{Q}(\Pi')$ be the compositum of $\mathbb{Q}(\mu)$ and $\mathbb{Q}(\Pi'_f)$. The following theorem, contained in Thm. 8.1, Thm. 8.6 and Prop. 8.7, generalizes the analogous results in the split case as well as Waldspurger's corresponding theorems for $d = 2$ and $m = 1$.

Theorem 1.5. *Let Π' be a regular algebraic representation of $G'(\mathbb{A})$ such that $JL(\Pi')$ is cuspidal. Then $\mathbb{Q}(\Pi')$ is a number field and Π'_f admits a $G'(\mathbb{A}_f)$-invariant $\mathbb{Q}(\Pi')$-structure. In particular, Π'_f is defined over a number field. Further, there is the equality of fields*

$$\mathbb{Q}(\Pi'_f) = \mathbb{Q}(JL(\Pi'_f)).$$

The above results on the nature of regular algebraic representations of $G'(\mathbb{A})$ suggest the following *schólion* to us.

Schólion. *Assuming the cuspidality of $JL(\Pi')$ – and not of Π' itself – is the appropriate setup to generalize statements on the arithmetic properties of cuspidal automorphic representations from GL_n/F to its inner forms.*

In view of this Schólion we may generalize Clozel's [8, Conjectures 3.7 and 3.8] as:

Conjecture 1.6. *Let Π' be a discrete series automorphic representation of $G'(\mathbb{A})$ such that $JL(\Pi')$ is cuspidal. Then the following are equivalent:*

(i) *Π'_f is defined over a number field.*
(ii) *Π' is algebraic.*

Beside the above theorems on the arithmetic properties of automorphic forms of $G'(\mathbb{A})$, we also prove in Sections 4 and 5 a number of purely representation-theoretical results on $GL_k(\mathbb{H})$ and $GL_n(\mathbb{R})$, $n = 2k$. To begin, we describe an explicit classification of the cohomological irreducible, unitary dual of $GL_k(\mathbb{H})$ and $GL_n(\mathbb{R})$, $n = 2k \geq 2$, following Vogan–Zuckerman [40]. This is contained in Thm. 4.9 and Thm. 4.12 which describe the set of all cohomological, irreducible, unitary $A_{\mathfrak{q}}(\lambda)$-modules of $GL_k(\mathbb{H})$ and $GL_n(\mathbb{R})$ with respect to any finite-dimensional coefficient system E_μ very concretely in terms of certain ordered partitions of k and n. The weight λ depends on the weight μ as in Definition 4.13 below. (This classification is also in accordance with the results of Speh, cf. [36].) Having parameterized the cohomological irreducible, unitary dual $\mathrm{Coh}_\mu(GL_k(\mathbb{H}))$ of $GL_k(\mathbb{H})$ and $\mathrm{Coh}_\mu(GL_n(\mathbb{R}))$ of $GL_n(\mathbb{R})$ with respect to E_μ by such partitions, we then prove the following theorem, cf. Thm. 5.2.

Theorem 1.7. *At a non-split archimedean place v of F, the local Jacquet-Langlands transfer $|LJ|_v$, constructed by Badulescu–Renard, defines a surjective map*

$$\mathrm{Coh}_\mu(GL_n(\mathbb{R})) \xrightarrow{|LJ|_v} \mathrel{\mkern-14mu}\twoheadrightarrow \mathrm{Coh}_\mu(GL_k(\mathbb{H}))$$

given explicitly by

$$|LJ|_v(A_{\mathfrak{q}_{\underline{n}}}(\lambda) \otimes \mathrm{sgn}^\varepsilon) = A_{\mathfrak{q}'_{\underline{k}}}(\lambda).$$

Here $\underline{n}$ (resp., $\underline{k}$) stands for the partition $n = \sum_{i=0}^r n_i$ (resp., $k = \sum_{i=0}^r k_i$) with $n_i = 2k_i$, $0 \leq i \leq r$ and $k_i > 0$ for $1 \leq i \leq r$.

Moreover, $|LJ|_v$ maps tempered cohomological representations to tempered cohomological representations.

Using the parametrization by ordered partitions, it is an easy exercise to determine the fibers of $|LJ|_v$ over a given $A_{\mathfrak{q}'}(\lambda)$-module. This theorem fits very well with the interplay of cohomological automorphic representations with Langlands functoriality as discussed in Raghuram–Shahidi in [28, Section 5.2].

Acknowledgements: A.R. thanks David Vogan for some email correspondence in 2003 concerning representations of $\mathrm{GL}_m(D)$ with cohomology which was the genesis of this project. However, at that time the generalized Jacquet-Langlands correspondence was still not proved and one had to wait for Badulescu's theorems. This project really got started when both the authors met at the Erwin Schrödinger Institute (ESI) in Vienna in February 2009. Both H.G. and A.R. thank ESI, and also the Max-Planck Institut für Mathematik for their hospitality during the time when the final version of this paper was written. H.G. also thanks the Department of Mathematics of the Oklahoma State University where much of this work was done.

2. The general linear group and its inner forms

2.1. **Generalities on division algebras.** Let F be a number field whose set of all places is denoted $V = V_\infty \cup V_f$, where as usual V_∞ is the subset of archimedean places and V_f the subset of non-archimedean places. The local completion of F at a place $v \in V$ is written F_v.

Let D be a central division-algebra over F of index d, i.e., $d^2 = \dim_F D$. The local algebras $D_v = D \otimes_F F_v$ are central simple algebras over F_v and hence isomorphic to a matrix algebra $M_{r_v}(A_v)$, for some integer $r_v \geq 1$ and a central division algebra A_v over F_v. The algebra D is said to be split at v if $A_v = F_v$ and non-split at v otherwise, i.e., A_v is not a field. The set of non-split places is finite. Analogous to the global situation, let d_v be the index of D_v, i.e., $d_v^2 = \dim_{F_v} A_v$. Then $r_v d_v = d$ for all v. If $v \in V_\infty$ is real then $d_v \in \{1, 2\}$, i.e., $A_v = \mathbb{R}$ or $\mathbb{H}$ and $D_v = M_d(\mathbb{R})$ if v is split and $M_{d/2}(\mathbb{H})$ is v is non-split (in which case d is even). Given any $m \geq 1$ we set $n := dm$ and $k := n/2$.

2.2. **The groups G' and G.** The determinant $\det'$ of an $m\times m$-matrix $X\in M_m(D)$, $m\geq 1$, is the generalization of the reduced norm to matrices: $\det'(X):=\det(\varphi(X\otimes 1))$, for some isomorphism $\varphi: M_m(D)\otimes_F\overline{\mathbb{Q}}\xrightarrow{\sim} M_n(\overline{\mathbb{Q}})$. It is independent of φ and is an F-rational polynomial in the coordinates of the entries of X. So the group

$$G'(F)=\{X\in M_m(D)|\det'(X)\neq 0\}$$

defines an algebraic group GL'_m over F. It is reductive and is an inner F-form of the split group $G:=GL_n/F$. At a real place $v\in V_\infty$ we hence obtain $G'(\mathbb{R})=GL_n(\mathbb{R})$ if v is split and $G'(\mathbb{R})=GL_k(\mathbb{H})$ if v is not split. We use the notation $G'_v:=G'(F_v)$ and $G_v:=G(F_v)$ for $v\in V$ and set as usual $G'_\infty:=\prod_{v\in V_\infty}G'_v$, resp., $G_\infty:=\prod_{v\in V_\infty}G_v$. Lie algebras of Lie groups are denoted by the same but gothic letter, e.g. $\mathfrak{g}_v=Lie(G_v)$, $\mathfrak{g}_\infty=Lie(G_\infty)$.

2.3. **Finite-dimensional representations.** We fix once and for all a maximal F-split torus T in G. The group $T_\infty=\prod_{v\in V_\infty}T_v=\prod_{v\in V_\infty}T(F_v)$ is then a Cartan subgroup of G_∞. Fixing the set of dominant algebraic characters $X^+(T_\infty)$ of T_∞ in the usual way, gives us that a tuple $\mu=(\mu_v)_{v\in V_\infty}\in X^+(T_\infty)$ can be identified with an equivalence class of irreducible, algebraic, finite-dimensional representations E_μ of G_∞ (on complex vector spaces) via the highest weight correspondence. It is clear that any such representation E_μ factors into irreducible representations $E_\mu=\bigotimes_{v\in V_\infty}E_{\mu_v}$, where E_{μ_v} is the irreducible representation of G_v of highest weight μ_v. If v is real, then $\mu_v=(\mu_{v,1},...,\mu_{v,n})$ with $\mu_{v,1}\geq...\geq\mu_{v,n}$ and $\mu_{v,i}\in\mathbb{Z}$, for $1\leq i\leq n$; and if v is complex, corresponding to the complex embeddings $\{\iota,\bar\iota\}$ of F, then μ_v is given by a pair $(\mu_{\iota_v},\mu_{\bar\iota_v})$ of n-tuples of the above form. A representation E_μ is called *essentially self-dual* if all its local factors E_{μ_v} are, i.e., if for all $v\in V_\infty$ there is a $w_v\in\mathbb{Z}$ such that $E_{\mu_v}\cong E^{\vee}_{\mu_v}\otimes\det^{w_v}$. At a real place this reads as

$$\mu_{v,i}+\mu_{v,n-i+1}=w_v,\qquad 1\leq i\leq n$$

and at a complex place this means

$$\mu_{\bar\iota_v,i}+\mu_{\iota_v,n-i+1}=w_v,\qquad 1\leq i\leq n.$$

It is called *self-dual* if $w_v=0$, i.e., $E_{\mu_v}\cong E^{\vee}_{\mu_v}$.

As G'_∞ is a real inner form of G_∞ the notion of highest weights and irreducible finite-dimensional representations is defined via the passage to the split form G_∞. That means that we say at a non-split place $v\in V_\infty$ an irreducible finite-dimensional representation E'_v of G'_v is *of highest weight* μ'_v, if the complexified representation E_v of $\mathfrak{g}'_v\otimes\mathbb{C}=\mathfrak{gl}_n(\mathbb{C})$ is. We hence drop the prime for such representations and write simply $E'_v=E_{\mu_v}$. Then, everything said above on representations of G_∞ also applies to irreducible finite-dimensional complex representations $E_\mu=\bigotimes_{v\in V_\infty}E_{\mu_v}$ of G'_∞ without changes (only adopting the notation of det to det$'$). A highest weight representation E_μ is called *regular*, if μ lies in the interior of the dominant Weyl chanber of G'_∞. The smallest algebraically integral element in the interior of the dominant Weyl chamber of G'_∞ and G_∞ is given by $\rho=(\rho_v)_{v\in V_\infty}$ with $\rho_v=(\frac{n-1}{2},\frac{n-3}{2},...,-\frac{n-1}{2})$ for all real places $v\in V_\infty$ (resp. the pair $\rho_v=(\rho_{\iota_v},\rho_{\bar\iota_v})$, $\rho_{\iota_v}=\rho_{\bar\iota_v}=(\frac{n-1}{2},\frac{n-3}{2},...,-\frac{n-1}{2})$, if v is complex).

2.4. **$(\mathfrak{g}'_\infty,K'^\circ_\infty)$-cohomology.** Let Z'/F be the center of the algebraic group G'/F and denote $Z'_\infty=\prod_{v\in V_\infty}Z'_v=\prod_{v\in V_\infty}Z'(F_v)$. At an archimedean place $v\in V_\infty$ we let K'_v be the product of a maximal compact subgroup of the real Lie group G'_v and Z'_v. Explicitly, we get

$$K'_v=\begin{cases} Sp(k)\mathbb{R}^* & \text{if } v \text{ non-split}\\ O(n)\mathbb{R}^* & \text{if } v \text{ split and real}\\ U(n)\mathbb{C}^* & \text{if } v \text{ complex,}\end{cases}$$

and define $K'_\infty=\prod_{v\in V_\infty}K'_v$. Analogously, we set $K_\infty=\prod_{v\in V_\infty}K_v$ where $K_v:=K'_v$ at split places and $K_v:=O(n)\mathbb{R}^*$ at non-split places. By K'°_∞ (resp., K°_∞) we denote the topological connected component of the identity within K'_∞ (resp., K_∞). Hence, locally

$$K'^\circ_v=\begin{cases} Sp(k)\mathbb{R}_+ & \text{if } v \text{ non-split}\\ SO(n)\mathbb{R}_+ & \text{if } v \text{ split and real}\\ U(n)\mathbb{C}^* & \text{if } v \text{ complex.}\end{cases}$$

We assume familiarity with the basic facts and notions concerning $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-modules (and $(\mathfrak{g}_\infty, K^\circ_\infty)$-modules), to be found in the book of Borel–Wallach [7], 0, I. All Lie-group representations $\Pi'_\infty = \bigotimes_{v\in V_\infty}\Pi'_v$ of G'_∞ appearing in this paper define a $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-module and hence $(\mathfrak{g}'_v, K'^\circ_v)$-modules, which we shall all denote by the same letter as the original Lie group representation. In particular, this applies to a highest weight representation $E_\mu = \bigotimes_{v\in V_\infty} E_{\mu_v}$. If furthermore, $\Pi'_\infty = \bigotimes_{v\in V_\infty}\Pi'_v$ is any $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-module, then we denote by

$$H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \Pi'_\infty)$$

its space of $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-cohomology (in degree q), cf. [7], I.5. A module Π'_∞ is called *cohomological*, if there is a highest weight representation E_μ as in Section 2.3, such that $H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \Pi'_\infty\otimes E_\mu)\neq 0$ for some degree q. It is a basic fact that these cohomology groups obey the Künneth-rule, i.e.,

$$H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \Pi'_\infty\otimes E_\mu) \cong \bigoplus_{\sum_v q_v = q}\ \bigotimes_{v\in V_\infty} H^{q_v}(\mathfrak{g}'_v, K'^\circ_v, \Pi'_v\otimes E_{\mu_v}).$$

Hence, Π'_∞ is cohomological, if and only if all its local components Π'_v are, i.e., they have non-vanishing $(\mathfrak{g}'_v, K'^\circ_v)$-cohomology with respect to some local highest weight representation E_{μ_v}.

3. Generalities on automorphic representations of $G'(\mathbb{A})$

3.1. We call an irreducible sub-quotient Π' of the space $\mathcal{A}(G'(F)\backslash G'(\mathbb{A}))$ of automorphic forms an *automorphic representation* of $G'(\mathbb{A})$ (although Π' is strictly speaking not a $G'(\mathbb{A})$-module), cf. the article of Borel–Jacquet [5] 3-4. Let $\mathbb{R}_+$ be the multiplicative group of positive real numbers, viewed as a subgroup

$$\mathbb{R}_+ \hookrightarrow G'(\mathbb{A})$$

by embedding it diagonally into G'_∞. The quotient $\mathbb{R}_+G'(F)\backslash G'(\mathbb{A})$ has finite volume (with respect to the usual quotient measure) and it therefore makes sense to talk about subspaces of square-integrable automorphic forms in $\mathcal{A}(\mathbb{R}_+G'(F)\backslash G'(\mathbb{A}))$. Now, recall that by its very definition, every automorphic form is annihilated by some power of an ideal $\mathcal{J}$ of finite codimension in the center of the universal enveloping algebra of $\mathfrak{g}'_\mathbb{C} = \mathfrak{g}'_\infty \otimes_\mathbb{R} \mathbb{C}$. Let us fix such an ideal $\mathcal{J}$ and denote by

$$\mathcal{A}_\mathcal{J}(G') \subset \mathcal{A}(\mathbb{R}_+G'(F)\backslash G'(\mathbb{A}))$$

the $G'(\mathbb{A})$-submodule consisting of those automorphic forms which are annihilated by some power of $\mathcal{J}$.

For later use, we will now recall a fine decomposition of the latter space $\mathcal{A}_\mathcal{J}(G')$, which was developed by Franke–Schwermer in [12] Thm. 1.4 and also in a similar way by Mœglin–Waldspurger in [27] III, Thm. 2.6., taking into account the so-called *parabolic support* and the *cuspidal support* of an automorphic representation. As a first step, $\mathcal{A}_\mathcal{J}(G')$ can be decomposed as a $G'(\mathbb{A})$-module into a finite direct sum, cf. [12] 1.1.(4),

$$\mathcal{A}_\mathcal{J}(G') \cong \bigoplus_{\{P'\}} \mathcal{A}_{\mathcal{J},\{P'\}}(G'), \tag{3.1}$$

ranging over the set of all *associate classes* $\{P'\}$ of a parabolic F-subgroup P' of G'. (Recall therefore that two parabolic F-subgroups P'_1 and P'_2 of G' are called associate, if their Levi-factors L'_1 and L'_2 are conjugate by an element in $G'(F)$.) More precisely, the spaces $\mathcal{A}_{\mathcal{J},\{P'\}}(G')$ consist exactly of those automorphic forms $f\in\mathcal{A}_\mathcal{J}(G')$, which are negligible along every parabolic F-subgroup $Q'\notin\{P'\}$, i.e., with respect to a Levi-decomposition of $Q' = L_{Q'}N_{Q'}$, the constant term $f_{Q'}$ is orthogonal to the space of cuspidal automorphic forms on $L_{Q'}(\mathbb{A})$.

We remark that within the direct sum (3.1), the subspace $\mathcal{A}_{cusp,\mathcal{J}}(G')$ of all cuspidal automorphic forms in $\mathcal{A}_\mathcal{J}(G')$ is given as the summand index by the class $\{G'\}$ itself:

$$\mathcal{A}_{cusp,\mathcal{J}}(G') = \mathcal{A}_{\mathcal{J},\{G'\}}(G').$$

3.2. The various summands $\mathcal{A}_{\mathcal{J},\{P'\}}(G')$ can be decomposed even further. Therefore, let $P' = L'N'$ be a Levi-decomposition of the parabolic F-subgroup P'. Its Levi factor L' is hence of the form $L' \cong \prod_{i=1}^{r} GL'_{m_i}$, with $\sum_{i=1}^{r} m_i = m$. Now, recall from [12], 1.2, the notion of an *associate class* $\varphi_{P'}$ of cuspidal automorphic representations of the Levi subgroups of the elements in the class $\{P'\}$. These classes $\varphi_{P'}$ may be parameterized by pairs of the form $(\chi, \tilde{\Pi}')$, where

1. $\tilde{\Pi}'$ is a unitary cuspidal automorphic representation of $L'(\mathbb{A})$, whose central character vanishes on the diagonally embedded group $\mathbb{R}^r_+ \hookrightarrow L'_\infty \hookrightarrow L'(\mathbb{A}) \cong \prod_{i=1}^{r} GL'_{m_i}(\mathbb{A})$
2. $\chi : \mathbb{R}^r_+ \to \mathbb{C}^*$ is a Lie group character and
3. the infinitesimal character of $\tilde{\Pi}'_\infty$ and the derivative $d\chi \in \mathrm{Hom}(\mathbb{C}^r, \mathbb{C})$ of χ are compatible with the action of $\mathcal{J}$ (cf. [12], 1.2).

Each associate class $\varphi_{P'}$ may hence be represented by a cuspidal automorphic representation $\Pi' := \tilde{\Pi}' \otimes e^{\langle d\chi, H_{P'}\rangle}$ of $L'(\mathbb{A})$, where $H_{P'}$ is the Harish-Chandra homomorphism $L'(\mathbb{A}) \to \mathbb{C}^r$, cf. [11], p. 185. Given $\varphi_{P'}$, represented by a cuspidal representation Π' of the above form, a $G'(\mathbb{A})$-submodule

$$\mathcal{A}_{\mathcal{J},\{P'\},\varphi_{P'}}(G')$$

of $\mathcal{A}_{\mathcal{J},\{P'\}}(G')$ was defined in [12], 1.3 as follows: It is the span of all possible holomorphic values or residues of all Eisenstein series attached to $\tilde{\Pi}'$, evaluated at the point $d\chi$, together with all their derivatives. This definition is independent of the choice of the representatives P' and Π', thanks to the functional equations satisfied by the Eisenstein series considered. For details, we refer the reader to [12] 1.2-1.4.

As a consequence of Franke's theorem, cf. [11], Thm. 14, the following refined decomposition as $G'(\mathbb{A})$-modules of the spaces $\mathcal{A}_{\mathcal{J},\{P'\}}(G')$ of automorphic forms was obtained in [12], Thm. 1.4:

$$\mathcal{A}_{\mathcal{J},\{P'\}}(G') \cong \bigoplus_{\varphi_{P'}} \mathcal{A}_{\mathcal{J},\{P'\},\varphi_{P'}}(G'). \tag{3.2}$$

This gives rise to the following

Definition 3.3. Let Ψ' be an automorphic representation of $G'(\mathbb{A})$, whose central character is trivial on the diagonally embedded group $\mathbb{R}_+$. If Ψ' is an irreducible subquotient of the space $\mathcal{A}_{\mathcal{J},\{P'\},\varphi_{P'}}(G')$, we call the associate class $\{P'\}$ a *parabolic support* and the associate class $\varphi_{P'}$ a *cuspidal support* of Ψ'.

3.3. **Discrete series representations.** Let Z'/F be the center of the algebraic group G'/F and

$$\omega : Z'(F)\backslash Z'(\mathbb{A}) \to \mathbb{C}^*$$

be a unitary, smooth character. The space

$$L^2_{dis}(G'(F)\backslash G'(\mathbb{A}), \omega)$$

denotes the space of all automorphic functions

$$f : G'(F)\backslash G'(\mathbb{A}) \to \mathbb{C}$$

which satisfy $f(zg) = \omega(z)f(g)$, for all $z \in Z'(\mathbb{A})$ and almost all $g \in G'(\mathbb{A})$, and $|f|^2$ is square-integrable as a function on $Z'(\mathbb{A})G'(F)\backslash G'(\mathbb{A})$ (with respect to the usual quotient measure). This space is the *discrete spectrum* of $G'(\mathbb{A})$, cf. Borel [3], 9.6. Via the right regular action, it is a representation space of $G'(\mathbb{A})$ and it decomposes as a direct Hilbert sum of irreducible, unitary $G'(\mathbb{A})$-representations $\tilde{\Pi}'$, called *unitary discrete series* representations:

$$L^2_{dis}(G'(F)\backslash G'(\mathbb{A}), \omega) = \widehat{\bigoplus} \tilde{\Pi}'.$$

Observe that there is no term for the multiplicity appearing in this decomposition, which is due to the Multiplicity One Theorem for unitary discrete series, proved by Badulescu–Renard in [2], Thm. 18.1.(b).

3.4. Let us now define a certain sub-class of automorphic representations, which we will mainly focus on in this paper.

Definition 3.4. Let $\mathscr{D}(G')$ be the family of all twists

$$\Pi' = \tilde{\Pi}' \otimes |\det{}'|^s,$$

$\tilde{\Pi}'$ being a unitary discrete series representation of $G'(\mathbb{A})$ and $s \in \mathbb{C}$. These twisted representations Π' are usually called *essentially discrete series* representations, but for brevity of terminology we will only say *discrete series* representations.

Clearly, $\mathscr{D}(G')$ contains the family of all twists of unitary cuspidal automorphic representations. In accordance with our previous terminology, we will henceforth call such a twist simply a *cuspidal* automorphic representation.

4. A CLASSIFICATION OF THE COHOMOLOGICAL, IRREDUCIBLE, UNITARY DUAL OF $GL_k(\mathbb{H})$ AND $GL_n(\mathbb{R})$

4.1. Being interested in arithmetic properties of automorphic forms of G'/F, the $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-cohomology of automorphic representations of $G'(\mathbb{A})$ will be an important tool. As we are planning to compare cohomological discrete series representations of the inner form $G'(\mathbb{A})$ to cohomological automorphic representations of $G(\mathbb{A})$, we shall in particular obtain some knowledge on the cohomological representations of the groups G'_v and G_v, v being a non-split archimedean place. In this section, we will hence determine the cohomological, irreducible, unitary dual of the real Lie groups G'_v and G_v, $v \in V_\infty$ non-split. This amounts in giving a classification of all irreducible unitary representations of $G'_v = GL_k(\mathbb{H})$, $k \geq 1$, and $G_v = GL_n(\mathbb{R})$, $n = 2k$, which have non-zero $(\mathfrak{g}'_v, K'^\circ_v)$-cohomology twisted by some irreducible highest weight representation E_{μ_v}.

Speh gave a list of irreducible unitary representations of $GL_n(\mathbb{R})$ in [36], which have non-zero cohomology with respect to the trivial representation $E_{\mu_v} = \mathbb{C}$. Using translation functors, see, e.g. [7] VI, sect. 0, this result can be adopted to the case of a general coefficient system E_{μ_v}. However, as we also want to classify the cohomological, irreducible, unitary dual of $GL_k(\mathbb{H})$, we are not going to use Speh's paper, but directly use the results of Vogan-Zuckerman, [40], in order to give a complete classification of the cohomological, irreducible, unitary dual of $GL_k(\mathbb{H})$ and $GL_n(\mathbb{R})$ in one go. To lighten the burden on the notation, we henceforth drag the subscript "v" about, keeping in mind that all objects are local ones at a non-split archimedean place.

4.2. **Non-equivalent θ'-stable parabolic subalgebras of $\mathfrak{g}'$.** Let θ' be the usual Cartan-involution $\theta'(X) = -\bar{X}^t$ on $\mathfrak{g}'$ ($\bar{X}$ denoting the standard conjugation of quaternionic matrices) giving rise to the Cartan decomposition $\mathfrak{g}' \cong \mathfrak{k}' \oplus \mathfrak{p}'$. According to this decomposition, let $\mathfrak{h}' = \mathfrak{t}' \oplus \mathfrak{a}'$ be a maximal compact, θ'-stable Cartan subalgebra. We take

$$\mathfrak{t}' = \left\{ \left(\begin{array}{ccc} ix_1 & & 0 \\ & \ddots & \\ 0 & & ix_k \end{array} \right) \middle| x_i \in \mathbb{R} \right\} \qquad \mathfrak{a}' = \left\{ \left(\begin{array}{ccc} y_1 & & 0 \\ & \ddots & \\ 0 & & y_k \end{array} \right) \middle| y_i \in \mathbb{R} \right\}.$$

For any $H \in \mathfrak{h}'_\mathbb{C}$, let $e_j(H) := ix_j$ and $f_j(H) := y_j$, $1 \leq j \leq k$. Then the set of roots of $\mathfrak{g}'_\mathbb{C}$ with respect to $\mathfrak{h}'_\mathbb{C}$ is given by

$$\Delta(\mathfrak{g}'_\mathbb{C}, \mathfrak{h}'_\mathbb{C}) = \{\pm e_i \pm e_j \pm (f_i - f_j), 1 \leq i < j \leq k\} \cup \{\pm 2e_i, 1 \leq i \leq k\}.$$

Furthermore,

$$\begin{array}{rcl} \Delta(\mathfrak{k}'_\mathbb{C}, \mathfrak{t}'_\mathbb{C}) & = & \{\pm e_i \pm e_j, 1 \leq i < j \leq k\} \cup \{\pm 2e_i, 1 \leq i \leq k\} \\ \Delta(\mathfrak{p}'_\mathbb{C}, \mathfrak{t}'_\mathbb{C}) & = & \{\pm e_i \pm e_j, 1 \leq i < j \leq k\} \\ W_{\mathfrak{k}'} & \cong & S_k \ltimes \{\pm 1\}^k, \end{array}$$

where S_k is the symmetric group of index k. It is hence clear that an element of the Weyl group $W_{\mathfrak{k}'}$ acts on $H \in i\mathfrak{t}'$ by permuting the entries x_j, $1 \leq j \leq k$ and changing their signs. This gives rise to the quotient $i\mathfrak{t}'/W_{\mathfrak{k}'}$, which admits a polyhedral cone decomposition. We obtain

Lemma 4.1. *The following sets are pairwise in* 1*-to-*1 *correspondence:*

(1) $\{$*open, polyhedral cones in* $i\mathfrak{t}'/W_{\mathfrak{k}'}\}$

(2) $\{x = (\underbrace{0,...,0}_{k_0},\underbrace{1,...,1}_{k_1},...,\underbrace{r,...,r}_{k_r}), k = \sum_{i=0}^{r} k_i, k_i > 0$ *for* $i \geq 1\}$

(3) $\{$*ordered partitions of* k*:* $\underline{k} = [k_0, k_1, ..., k_r], k_0 \geq 0, k_i > 0$ *for* $i \geq 1\}$

Proof. The existence of the bijection (1)↔(2) is a direct consequence of the description of the action of $W_{\mathfrak{k}'}$ on $i\mathfrak{t}'$. (The j-th entry of x is its coordinate with respect to the basis element of $i\mathfrak{t}'$, which is dual to e_{k-j+1}.) The bijection (2)↔(3) holds trivially. □

Now, let $\mathfrak{Q}'$ be the set of all θ'-stable parabolic subalgebras $\mathfrak{q}'$ of $\mathfrak{g}'_{\mathbb{C}}$, which contain a fixed Borel subalgebra. Recall that $K' \cong Sp(k)\mathbb{R}^*$ acts on this set by the adjoint action. This leads to a finite set $\mathfrak{Q}'/K'$ of orbits or - otherwise put - K'-conjugacy classes of θ'-stable parabolic subalgebras $\mathfrak{q}'$ containing a fixed Borel subalgebra. The following lemma says that the set of open polyhedral cones in $i\mathfrak{t}'/W_{\mathfrak{k}'}$ and $\mathfrak{Q}'/K'$ are in bijection.

Lemma 4.2 ([21] IV, Prop. 4.76)**.** *Every* $x \in i\mathfrak{t}'/W_{\mathfrak{k}'}$ *defines a* θ'*-stable, parabolic subalgebra* $\mathfrak{q}'_x$ *in* $\mathfrak{g}'_{\mathbb{C}}$ *by assigning it a Levi decomposition* $\mathfrak{q}'_x := \mathfrak{l}'_x \oplus \mathfrak{u}'_x$*, where*

$$\mathfrak{l}'_x := \mathfrak{h}'_{\mathbb{C}} \oplus \bigoplus_{\substack{\alpha\in\Delta(\mathfrak{g}'_{\mathbb{C}},\mathfrak{t}'_{\mathbb{C}})\\ \alpha(x)=0}} (\mathfrak{g}'_{\mathbb{C}})_\alpha \qquad \text{and} \qquad \mathfrak{u}'_x := \bigoplus_{\substack{\alpha\in\Delta(\mathfrak{g}'_{\mathbb{C}},\mathfrak{t}'_{\mathbb{C}})\\ \alpha(x)>0}} (\mathfrak{g}'_{\mathbb{C}})_\alpha$$

and two such x_1*,* x_2 *define the same parabolic subalgebra* $\mathfrak{q}'_{x_1} = \mathfrak{q}'_{x_2}$ *if and only if they are in the same open, polyhedral cone.*
Conversely, up to conjugacy by K'*, any* θ'*-stable, parabolic subalgebra* $\mathfrak{q}'$ *containing a fixed Borel subalgebra is of the form* $\mathfrak{q} = \mathfrak{q}_x$*, with* $x \in i\mathfrak{t}'/W_{\mathfrak{k}'}$*.*

Together with Lem. 4.1, the last lemma provides a parametrization of the set of K'-conjugacy classes of θ'-stable parabolic subalgebras $\mathfrak{q}'$ of $\mathfrak{g}'_{\mathbb{C}}$ containing a fixed Borel subalgebra by ordered partitions $\underline{k} = [k_0, k_1, ..., k_r]$ of k: $k = \sum_{i=0}^{r} k_i$, $k_0 \geq 0$ and $k_i > 0$ for $i \geq 1$. Therefore, we shall henceforth write $\mathfrak{q}'_{\underline{k}}$ for a class in $\mathfrak{Q}'/K'$. Its Levi subalgebra has real part

$$\mathfrak{l}'^{\mathbb{R}}_{\underline{k}} := \mathfrak{l}'_{\underline{k}} \cap \mathfrak{g}' \cong \mathfrak{gl}_{k_0}(\mathbb{H}) \oplus \bigoplus_{i=1}^{r} \mathfrak{gl}_{k_i}(\mathbb{C}). \tag{4.3}$$

In view of Salamanca-Riba's paper [32], Prop. 1.11, we shall introduce another relation on $\mathfrak{Q}'/K'$: $\mathfrak{q}'_{\underline{k}_1} \sim \mathfrak{q}'_{\underline{k}_2}$ if and only if $\Delta(\mathfrak{u}'_{\underline{k}_1} \cap \mathfrak{p}'_{\mathbb{C}}, \mathfrak{t}'_{\mathbb{C}}) = \Delta(\mathfrak{u}'_{\underline{k}_2} \cap \mathfrak{p}'_{\mathbb{C}}, \mathfrak{t}'_{\mathbb{C}})$. It is an easy exercise to check that in terms of the parameterizing partitions $\underline{k}_1$ and $\underline{k}_2$, $\mathfrak{q}'_{\underline{k}_1} \sim \mathfrak{q}'_{\underline{k}_2}$ if and only if either $\underline{k}_1 = \underline{k}_2$ or

$$\underline{k}_1 = [0, 1, k_2, ..., k_r] \qquad \text{and} \qquad \underline{k}_2 = [1, k_2, ..., k_r]. \tag{4.4}$$

(E.g., $\underline{k}_1 = [0, 1, 2, 1, 3]$ and $\underline{k}_2 = [1, 2, 1, 3]$ for $k = 7$.) Within such a non-singleton equivalence class, we pick the K'-conjugacy class parameterized by $\underline{k}_1$. To conclude this subsection, we define $\mathcal{Q}'$ to be the so chosen set of representatives of equivalence classes in $\mathfrak{Q}'/K'$, i.e., of K'-conjugacy classes of θ'-stable parabolic subalgebras $\mathfrak{q}' = \mathfrak{q}'_{\underline{k}}$ of $\mathfrak{g}'_{\mathbb{C}}$ containing a fixed Borel subalgebra.

4.3. **Non-equivalent θ-stable parabolic subalgebras of $\mathfrak{g}$.** We will now determine the same data for the split case. Therefore, let θ be the usual Cartan-involution $\theta(X) = -X^t$ on $\mathfrak{g}$ leading to the Cartan decomposition $\mathfrak{g} \cong \mathfrak{k} \oplus \mathfrak{p}$ and standard maximal compact, θ-stable Cartan subalgebra $\mathfrak{h} = \mathfrak{t} \oplus \mathfrak{a}$. We have

$$\mathfrak{t} = \left\{ \begin{pmatrix} 0 & x_1 & & & 0 \\ -x_1 & 0 & & & \\ & & \ddots & & \\ & & & 0 & x_k \\ & & & -x_k & 0 \end{pmatrix} \middle| x_i \in \mathbb{R} \right\}$$

and

$$\mathfrak{a} = \left\{ \left(\begin{array}{ccccc} y_1 & & & & \\ & y_1 & & & \\ & & \ddots & & \\ & & & y_k & \\ & & & & y_k \end{array} \right) \middle| y_i \in \mathbb{R} \right\}.$$

For $H \in \mathfrak{h}_{\mathbb{C}}$, let again $e_j(H) := ix_j$ and $f_j(H) := y_j$, $1 \leq j \leq k$. Then the set of roots of $\mathfrak{g}_{\mathbb{C}}$ with respect to $\mathfrak{h}_{\mathbb{C}}$ is given by

$$\Delta(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) = \{\pm e_i \pm e_j \pm (f_i - f_j), 1 \leq i < j \leq k\} \cup \{\pm 2e_i, 1 \leq i \leq k\}.$$

However, in contrast to the non-split case, the sets of compact and non-compact roots change their roles

$$\begin{aligned} \Delta(\mathfrak{k}_{\mathbb{C}}, \mathfrak{t}_{\mathbb{C}}) &= \{\pm e_i \pm e_j, 1 \leq i < j \leq k\} \\ \Delta(\mathfrak{p}_{\mathbb{C}}, \mathfrak{t}_{\mathbb{C}}) &= \{\pm e_i \pm e_j, 1 \leq i < j \leq k\} \cup \{\pm 2e_i, 1 \leq i \leq k\}. \end{aligned}$$

Slightly abusing notation, we set

$$W_{\mathfrak{k}} := S_k \ltimes \{\pm 1\}^k,$$

and let this "extended" Weyl group $W_{\mathfrak{k}}$ act on $H \in i\mathfrak{t}$ by permuting the entries x_j, $1 \leq j \leq k$ and changing their signs. Whence, the following lemma is obvious by Lem. 4.1. See also Speh's aforementioned article [36], p. 464.

Lemma 4.5. *The following sets are pairwise in 1-to-1 correspondence:*

(1) $\{$*open, polyhedral cones in* $i\mathfrak{t}/W_{\mathfrak{k}}\}$
(2) $\{x = (\underbrace{0,...,0}_{k_0}, \underbrace{1,...,1}_{k_1}, ..., \underbrace{r,...,r}_{k_r}), k = \sum_{i=0}^{r} k_i, k_i > 0$ *for* $i \geq 1\}$
(3) $\{$*ordered partitions of* n*:* $\underline{n} = [n_0, n_1, ..., n_r], n_0 \geq 0, n_i = 2k_i$*, where* $k_i > 0$ *for* $i \geq 1\}$

Now, let $\mathfrak{Q}$ be the set of all θ-stable parabolic subalgebras $\mathfrak{q}$ of $\mathfrak{g}_{\mathbb{C}}$, which contain a fixed Borel subalgebra. Recall that $K \cong O(n)\mathbb{R}^*$ acts on this set by the adjoint action. This leads a finite set $\mathfrak{Q}/K$ of K-conjugacy classes of θ-stable parabolic subalgebras $\mathfrak{q}$ containing a fixed Borel subalgebra. Again there is a bijection between the set of open polyhedral cones in $i\mathfrak{t}/W_{\mathfrak{k}}$ and $\mathfrak{Q}/K$, explaining our choice of the group $W_{\mathfrak{k}}$:

Lemma 4.6 ([21] IV, Prop. 4.76)**.** *Every* $x \in i\mathfrak{t}/W_{\mathfrak{k}}$ *defines a* θ*-stable, parabolic subalgebra* $\mathfrak{q}_x$ *in* $\mathfrak{g}_{\mathbb{C}}$ *via* $\mathfrak{q}_x := \mathfrak{l}_x \oplus \mathfrak{u}_x$*, where*

$$\mathfrak{l}_x := \mathfrak{h}_{\mathbb{C}} \oplus \bigoplus_{\substack{\alpha \in \Delta(\mathfrak{g}_{\mathbb{C}}, \mathfrak{t}_{\mathbb{C}}) \\ \alpha(x) = 0}} (\mathfrak{g}_{\mathbb{C}})_\alpha \qquad \text{and} \qquad \mathfrak{u}_x := \bigoplus_{\substack{\alpha \in \Delta(\mathfrak{g}_{\mathbb{C}}, \mathfrak{t}_{\mathbb{C}}) \\ \alpha(x) > 0}} (\mathfrak{g}_{\mathbb{C}})_\alpha$$

and two such x_1*,* x_2 *define the same parabolic subalgebra* $\mathfrak{q}_{x_1} = \mathfrak{q}_{x_2}$ *if and only if they are in the same open, polyhedral cone.*
Conversely, up to conjugacy by K*, any* θ*-stable, parabolic subalgebra* $\mathfrak{q}$ *containing a fixed Borel subalgebra of* $\mathfrak{g}_{\mathbb{C}}$ *is of the form* $\mathfrak{q} = \mathfrak{q}_x$*, with* $x \in i\mathfrak{t}/W_{\mathfrak{k}}$*.*

In particular, Lem. 4.5 provides a parametrization of the set of K-conjugacy classes of θ-stable parabolic subalgebras $\mathfrak{q}$ of $\mathfrak{g}_{\mathbb{C}}$ containing a fixed Borel subalgebra by ordered partitions $\underline{n} = [n_0, n_1, ..., n_r]$ of n: $n = \sum_{i=0}^{r} n_i$, $n_0 \geq 0$ and $n_i = 2k_i > 0$ for $i \geq 1$. Therefore, we shall henceforth write $\mathfrak{q}_{\underline{n}}$ for a class in $\mathfrak{Q}/K$. The real part of its Levi subalgebra is isomorphic to

$$\mathfrak{l}^{\mathbb{R}}_{\underline{n}} := \mathfrak{l}_{\underline{n}} \cap \mathfrak{g} \cong \mathfrak{gl}_{n_0}(\mathbb{R}) \oplus \bigoplus_{i=1}^{r} \mathfrak{gl}_{k_i}(\mathbb{C}), \tag{4.7}$$

see also [36], p. 464. Introducing the analogous equivalence relation on $\mathfrak{Q}/K$ by letting $\mathfrak{q}_{\underline{n}_1} \sim \mathfrak{q}_{\underline{n}_2}$ if and only if $\Delta(\mathfrak{u}_{\underline{n}_1} \cap \mathfrak{p}_{\mathbb{C}}, \mathfrak{t}_{\mathbb{C}}) = \Delta(\mathfrak{u}_{\underline{n}_2} \cap \mathfrak{p}_{\mathbb{C}}, \mathfrak{t}_{\mathbb{C}})$, we encounter a different phenomenon than in the non-split case.

As every root appears at least once in the set of non-compact roots, for any θ-stable parabolic subalgebra $\mathfrak{q}_{\underline{n}}$ of $\mathfrak{g}_{\mathbb{C}}$

$$\Delta(\mathfrak{u}_{\underline{n}} \cap \mathfrak{p}_{\mathbb{C}}, \mathfrak{t}_{\mathbb{C}}) = \Delta(\mathfrak{u}_{\underline{n}}, \mathfrak{t}_{\mathbb{C}})$$

and so the relation "$\sim$" degenerates to equality:

$$\mathfrak{q}_{\underline{n}_1} \sim \mathfrak{q}_{\underline{n}_2} \quad \Leftrightarrow \quad \mathfrak{q}_{\underline{n}_1} = \mathfrak{q}_{\underline{n}_2}.$$

Again, we denote by $\mathcal{Q}$ the set of all such equivalence classes, i.e., $\mathcal{Q} = \mathfrak{Q}/K$ itself in this case.

4.4. Having compiled this data for $\mathfrak{g}'$ and $\mathfrak{g}$, we can now give the desired classification of the cohomological, irreducible, unitary dual of G' and G. Therefore, let E be an irreducible representation of G' or G on a finite-dimensional complex vector space. Observe that we have changed Cartan subalgebras in this section, so we will write λ for its highest weight with respect to the standard choice of positivity on $\Delta(\mathfrak{g}_{\mathbb{C}}, \mathfrak{h}_{\mathbb{C}}) = \Delta(\mathfrak{g}'_{\mathbb{C}}, \mathfrak{h}'_{\mathbb{C}})$, rather than μ. As we will soon use the results of Vogan–Zuckerman [40], Section 5, we shall check when λ does define a so-called *admissible* character: Given $\mathfrak{q}' \in \mathcal{Q}'$ (resp., $\mathfrak{q} \in \mathcal{Q}$) let $L' \subseteq G'$ (resp., $L \subseteq G$) be the connected subgroup with Lie algebra $\mathfrak{l}'^{\mathbb{R}}$ (resp., $\mathfrak{l}^{\mathbb{R}}$). According to [40], (5.1), a linear functional ξ on $\mathfrak{l}'$ (resp., $\mathfrak{l}$) is called admissible if

(1) ξ is the differential of a unitary character of L' (resp., L)
(2) $\langle \xi, \alpha \rangle \geq 0$ for all roots α appearing in $\mathfrak{u}'$ (resp., $\mathfrak{u}$).

As a corollary, a highest weight λ defines an admissible character of $\mathfrak{l}'$ if and only if

$$\lambda|_{[\mathfrak{l}'^{\mathbb{R}}, \mathfrak{l}'^{\mathbb{R}}]} = 0 \quad \text{and} \quad \lambda|_{\mathfrak{a}'} = 0. \tag{4.8}$$

As an explanation, the vanishing of λ on the commutator $[\mathfrak{l}'^{\mathbb{R}}, \mathfrak{l}'^{\mathbb{R}}]$ is equivalent to saying that $\lambda : \mathfrak{h}'_{\mathbb{C}} \to \mathbb{C}$ can be extended to a character on $\mathfrak{l}' \supseteq \mathfrak{h}'_{\mathbb{C}}$, while the second condition $\lambda|_{\mathfrak{a}'} = 0$ is equivalent to λ coming from a *unitary* character. The latter condition can once more be equivalently reformulated by saying that $\lambda \circ \theta' = \lambda$ or - again equivalent - that E is self-dual. In particular, we may view λ as being expressed in the functionals e_j, $1 \leq j \leq k$, only, writing $\lambda = (\lambda_1, ..., \lambda_k)$, λ_j being the coefficient of e_j in such a decomposition. Clearly, the same statements hold for $\mathfrak{g}'$ being replaced by $\mathfrak{g}$, removing the prime everywhere.

We let $\mathcal{Q}'(\lambda)$ (resp., $\mathcal{Q}(\lambda)$) be the set of all $\mathfrak{q}' \in \mathcal{Q}'$ (resp., $\mathfrak{q} \in \mathcal{Q}$), to whose Levi parts $\mathfrak{l}'$ (resp., $\mathfrak{l}$) a given highest weight λ can be admissibly extended.

4.5. **The cohomological, irreducible, unitary dual of $G' = GL_k(\mathbb{H})$.** Now, we may state the following theorem settling the non-split case.

Theorem 4.9. *Let $G' = GL_k(\mathbb{H})$, $k \geq 1$ and $E = E_\lambda$ a highest weight representation. The following holds:*

(1) *To each $\mathfrak{q}' = \mathfrak{q}'_{\underline{k}} \in \mathcal{Q}'(\lambda)$, there exists a unique irreducible unitary G'-representation $A_{\mathfrak{q}'}(\lambda)$. None of these are pairwise isomorphic, i.e., $A_{\mathfrak{q}'_{\underline{k}_1}}(\lambda) \cong A_{\mathfrak{q}'_{\underline{k}_2}}(\lambda)$ if and only if $\mathfrak{q}'_{\underline{k}_1} = \mathfrak{q}'_{\underline{k}_2}$.*
(2) *Each $A_{\mathfrak{q}'}(\lambda)$ is cohomological with respect to E_λ, i.e.,*

$$H^q(\mathfrak{g}', K'^\circ, A_{\mathfrak{q}'}(\lambda) \otimes E_\lambda) \neq 0$$

for some degree $q \geq 0$, and all irreducible unitary G'-modules which are cohomological with respect to E_λ are obtained in this way.
(3) *The Poincaré-polynomial of the cohomology ring $H^*(\mathfrak{g}', K'^\circ, A_{\mathfrak{q}'(\lambda)} \otimes E_\lambda)$ is independent of λ and given by*

$$P(\underline{k}, X) = \frac{X^{\dim(\mathfrak{u}'_{\underline{k}} \cap \mathfrak{p}'_{\mathbb{C}})}}{1+X} \prod_{i=1}^{r} \prod_{j=1}^{k_i} (1 + X^{2j-1}) \prod_{j=1}^{k_0} (1 + X^{4j-3}).$$

(4) *Among all $A_{\mathfrak{q}'}(\lambda)$, $\mathfrak{q}' = \mathfrak{q}'_{\underline{k}} \in \mathcal{Q}'(\lambda)$, the representation indexed by $\mathfrak{q}'_{\underline{k}}$ with $\underline{k} = [0, 1, 1, ..., 1]$ is the only tempered representation.*

Proof. (1): The existence of the irreducible G'-module $A_{\mathfrak{q}'}(\lambda)$ is proved by Vogan–Zuckerman in [40], Thm. 5.3 together with (5.1). Its unitarity is shown by Vogan in [39], Thm. 1.3, while their pairwise inequivalence is a consequence of the work of Salamanca-Riba [32], Prop. 1.11.
(2): This is [40], Thm. 5.5 and 5.6.
(3): By [40], Thm. 5.5 we know that

$$H^q(\mathfrak{g}', K'^\circ, A_{\mathfrak{q}'_{\underline{k}}}(\lambda) \otimes E_\lambda) \cong H^{q-\dim(\mathfrak{u}'_{\underline{k}}\cap\mathfrak{p}'_{\mathbb{C}})}(\mathfrak{l}'^{\mathbb{R}}_{\underline{k}}, \mathfrak{l}'^{\mathbb{R}}_{\underline{k}} \cap \mathfrak{k}', \mathbb{C}),$$

which is already independent of λ but only dependent on the partition $\underline{k}$ determining $\mathfrak{q}' = \mathfrak{q}'_{\underline{k}}$. Using the concrete form of $\mathfrak{l}'^{\mathbb{R}}_{\underline{k}}$, given in (4.3) and the fact that $(\mathfrak{g}', K'^\circ)$-cohomology satisfies the Künneth-rule, it is now an easy exercise to calculate the Poincaré-polynomial of the cohomology ring $H^*(\mathfrak{g}', K'^\circ, A_{\mathfrak{q}'}(\lambda) \otimes E_\lambda)$.
(4): Taking into account our description of the real part of the Levi subalgebra of a θ'-stable parabolic $\mathfrak{q}'_{\underline{k}} \in \mathcal{Q}'(\lambda)$ given in (4.3), this follows from [40], last paragraph on p. 58. ☐

Remark 4.10. It is to avoid redundancies in the list of $A_{\mathfrak{q}'}(\lambda)$-modules, that we introduced the equivalence relation "$\sim$" on $\mathfrak{Q}'/K'$. If there is a non-trivial equivalence $\mathfrak{q}'_{\underline{k}_1} \sim \mathfrak{q}'_{\underline{k}_2}$, i.e., $\underline{k}_1 = [0, 1, k_2, ..., k_r]$ and $\underline{k}_2 = [1, k_2, ..., k_r]$, and λ can be extended to an admissible character of $\mathfrak{l}'_{\underline{k}_2}$, too, then $A_{\mathfrak{q}'_{\underline{k}_2}}(\lambda)$ exists. However, it such a case

$$A_{\mathfrak{q}'_{\underline{k}_1}}(\lambda) \cong A_{\mathfrak{q}'_{\underline{k}_2}}(\lambda),$$

cf. [32], Prop. 1.11.

Thm. 4.9 motivates the following definition.

Definition 4.11. Let $E = E_\mu$ be a highest weight representation as in Section 2.3, whose highest weight with respect to $\mathfrak{h}'_{\mathbb{C}}$ is λ. We denote

$$\mathrm{Coh}_\mu(G') := \{A_{\mathfrak{q}'}(\lambda), \mathfrak{q}' = \mathfrak{q}'_{\underline{k}} \in \mathcal{Q}'(\lambda)\}.$$

According to Thm. 4.9, $\mathrm{Coh}_\mu(G')$ is the set of all irreducible, unitary G'-representations, which have non-zero $(\mathfrak{g}', K'^\circ)$-cohomology with respect to E.

4.6. The cohomological, irreducible, unitary dual of $G = GL_n(\mathbb{R})$.

The split case is slightly more complicated, as $G = GL_n(\mathbb{R})$ is not connected. Still, we obtain the following theorem.

Theorem 4.12. *Let $G = GL_n(\mathbb{R})$, $n = 2k \geq 2$ and $E = E_\lambda$ a highest weight representation. The following holds:*

1. *To each $\mathfrak{q} = \mathfrak{q}_{\underline{n}} \in \mathcal{Q}(\lambda)$, there exists a unique irreducible unitary G-representation $A_{\mathfrak{q}}(\lambda)$. None of these are pairwise isomorphic, i.e., $A_{\mathfrak{q}_{\underline{n}_1}}(\lambda) \cong A_{\mathfrak{q}_{\underline{n}_2}}(\lambda)$ if and only if $\mathfrak{q}_{\underline{n}_1} = \mathfrak{q}_{\underline{n}_2}$.*
2. *Each $A_{\mathfrak{q}}(\lambda)$ is cohomological with respect to E_λ, i.e.,*
$$H^q(\mathfrak{g}, K^\circ, A_{\mathfrak{q}}(\lambda) \otimes E_\lambda) \neq 0$$
for some degree $q \geq 0$. Conversely, all irreducible unitary G-modules which are cohomological with respect to E_λ are of the form
$$A_{\mathfrak{q}}(\lambda) \otimes \mathrm{sgn}^\varepsilon, \qquad \mathfrak{q} = \mathfrak{q}_{\underline{n}} \in \mathcal{Q}(\lambda), \varepsilon \in \{0, 1\}.$$
3. *Among all $A_{\mathfrak{q}}(\lambda)$, $\mathfrak{q} = \mathfrak{q}_{\underline{n}} \in \mathcal{Q}(\lambda)$, the representation indexed by $\mathfrak{q}_{\underline{n}}$ with $\underline{n} = [0, 2, 2, ..., 2]$ is the only tempered representation.*

Proof. (1): The existence of the irreducible G'-module $A_{\mathfrak{q}'}(\lambda)$ is again a consequence [40], Thm. 5.3 together with (5.1). Its unitarity is shown in [39], Thm. 1.3, while their pairwise inequivalence is a consequence of [32], Prop. 1.11.
(2): This is [40], Thm. 5.5 and 5.6., where the twist by sgn^ε takes account of the disconnectedness of G.
(3): Recalling our concrete description of the real part of the Levi subalgebra of a θ-stable parabolic $\mathfrak{q}_{\underline{n}} \in \mathcal{Q}(\lambda)$ given in (4.7), this follows again from [40], last paragraph on p. 58. ☐

Definition 4.13. Let $E = E_\mu$ be a highest weight representation as in Section 2.3, whose highest weight with respect to $\mathfrak{h}_\mathbb{C}$ is λ. We denote

$$\mathrm{Coh}_\mu(G) := \{A_\mathfrak{q}(\lambda) \otimes \mathrm{sgn}^\varepsilon, \mathfrak{q} = \mathfrak{q}_{\underline{n}} \in \mathcal{Q}(\lambda), \varepsilon \in \{0,1\}\}.$$

According to Thm. 4.12, $\mathrm{Coh}_\mu(G)$ is the set of all irreducible, unitary G-representations, which have non-zero $(\mathfrak{g}, K^\circ)$-cohomology with respect to E.

5. The local and global Jacquet-Langlands transfer and its interplay with cohomology

5.1. **The local Jacquet-Langlands map.** In [20] Jacquet–Langlands established a bijection between irreducible representations of G'_v and irreducible, admissible square-integrable representations of G_v in the special case when $m = 1$ and $d = 2$, i.e., when G' is the group of invertible elements in a quaternion division algebra. (See also Gelbart–Jacquet [13] Thm. 8.1 and its generalization of Rogwaski, [30].) This bijection was extended to a bijection between irreducible, admissible square-integrable representations of G'_v and irreducible, admissible square-integrable representations of G_v for any m and d by Deligne–Kazhdan–Vignéras in [10] – the local Jacquet-Langlands correspondence. Finally, Badulescu in [1], p. 406 together with Badulescu–Renard in [2], sect. 13, “thickened” the Jacquet-Langlands correspondence to a map

$$|LJ|_v : \tilde{U}_{cp}(G_v) \to \tilde{U}(G'_v)$$

from the family $\tilde{U}_{cp}(G_v)$ of so-called d_v*-compatible* irreducible unitary representations of G_v to the family $\tilde{U}(G'_v)$ of irreducible unitary representations of G'_v. (For the notion of d_v-compatibility we refer to [1], sect. 15) This map is neither injective nor surjective in general, but restricts to the Jacquet-Langlands correspondence of [10] (and [20]) on unitary square-integrable representations. If $d_v = 1$, i.e., $G'_v = G_v$ the map $|LJ|_v$ is the identity. Furthermore, $|LJ|_v$ commutes with parabolic induction and forming tensor products. We denote by $U_{cp}(G_v)$ the set of all $\Pi_v = \tilde{\Pi}_v \otimes |\det|^s$, with $s \in \mathbb{C}$ and $\tilde{\Pi}_v \in \tilde{U}_{cp}(G_v)$, and define

$$|LJ|_v(\Pi_v) = |LJ|_v(\tilde{\Pi}_v) \otimes \det'^s.$$

The analogous notation $U(G'_v)$ is used in the non-split case, i.e., $\Pi'_v \in U(G'_v)$ if and only if $\Pi'_v = \tilde{\Pi}'_v \otimes |\det'|^s$ with $\tilde{\Pi}'_v \in \tilde{U}(G'_v)$. Furthermore we set for $\Pi_\infty = \bigotimes_{v \in V_\infty} \Pi_v \in U_{cp}(G_\infty) = \bigotimes_{v \in V_\infty} U_{cp}(G_v)$

$$|LJ|_\infty(\Pi_\infty) = \bigotimes_{v \in V_\infty} |LJ|_v(\Pi_v).$$

5.2. In the sequel, we will need some particular knowledge about the easiest, non-trivial case of $|LJ|_v$, namely if $v \in V_\infty$ is a non-split place and $m = 1$ and $d = 2$. For any integer $l \geq 1$ and $u \in \mathbb{C}$ denote by

$$F(u, l) := \mathrm{Sym}^{l-1}\mathbb{C}^2 \otimes \det'^{-u/2},$$

where $\mathrm{Sym}^{l-1}\mathbb{C}^2$ is the unique irreducible, unitary representation of $SL_1(\mathbb{H})$ of dimension l. The representations $F(u, l)$ exhaust all irreducible, finite-dimensional representations of $GL_1(\mathbb{H})$.

Furthermore, we define

$$D(u, l) := D(l) \otimes |\det|^{-u/2},$$

where $D(l)$ the unique irreducible, unitary, discrete series representation of $SL_2^\pm(\mathbb{R})$ of lowest (non-negative) $O(2)$-type $l + 1$. Then we obtain

Lemma 5.1. *Let $v \in V_\infty$ be a non-split place of a quaternion division algebra D and $m = 1$. The family $U_{cp}(GL_2(\mathbb{R}))$ consists precisely of the representations*

- (i) $\mathrm{sgn}^\varepsilon |\det|^s$, $\varepsilon \in \{0, 1\}$ *and* $s \in \mathbb{C}$
- (ii) $D(u, l)$ *for some integer* $l \geq 1$ *and* $u \in \mathbb{C}$.

while the family $U(GL_1(\mathbb{H}))$ consists precisely of the finite-dimensional representations

- (i') $F(u', l')$ *for some integer* $l' \geq 1$ *and* $u' \in \mathbb{C}$.

The local map $|LJ|_v$ *is given by*

$$|LJ|_v(\Pi_v) = \begin{cases} \det'^{s} & \textit{if } \Pi_v = \mathrm{sgn}^{\varepsilon}|\det|^{s} \\ F(u,l) & \textit{if } \Pi_v = D(u,l). \end{cases}$$

In particular, in this special case, $|LJ|_v$ *is surjective but not injective.*

Proof. This is well-known, respectively follows from the description of the local map $|LJ|_v$ in this case, cf. Badulescu–Renard [2], Thm 13.8. ☐

5.3. **Local Jacquet-Langlands and cohomology.** We are now ready to prove the first main result of this paper. It compares the cohomological, irreducible, unitary duals of G'_v and G_v, $v \in V_\infty$ non-split, via the local Jacquet-Langlands map $|LJ|_v$ and will be of particular importance in the following sections. Nevertheless, we believe that it is interesting in its own right. It fits very well with the philosophy to use functoriality in order to get cohomological automorphic representations. The interested reader may find a survey on this topic by Raghuram–Shahidi in [28], Section 5.2.

Theorem 5.2. *Let* $v \in V_\infty$ *be a non-split place, so* $G'_v = GL_k(\mathbb{H})$, $k \geq 1$ *and* $G_v = GL_n(\mathbb{R})$, $n = 2k$. *Furthermore, let* $E = E_{\mu_v}$ *be any highest weight representation as in Section 2.3. Then we get:*
The local Jacquet-Langlands map $|LJ|_v$ *defines a surjection*

$$\begin{tikzcd} \mathrm{Coh}_{\mu_v}(G_v) \arrow[r, two heads, "|LJ|_v"] & \mathrm{Coh}_{\mu_v}(G'_v) \end{tikzcd}$$

given explicitly by

$$|LJ|_v(A_{\mathfrak{q}_{\underline{n}}}(\lambda) \otimes \mathrm{sgn}^{\varepsilon}) = A_{\mathfrak{q}'_{\underline{k}}}(\lambda).$$

Here $\underline{n} = [n_0, n_1, ..., n_r]$ *and* $\underline{k} = [k_0, k_1, ..., k_r]$ *with* $n_i = 2k_i$, $0 \leq i \leq r$.
In particular, $|LJ|_v$ *maps tempered cohomological representations to tempered cohomological representations.*

Proof. The proof consists in rewriting the $A_{\mathfrak{q}'}(\lambda)$- and $A_{\mathfrak{q}}(\lambda)$-modules as quotients of parabolically induced representations, following Vogan–Zuckerman [40], p. 82, and then applying $|LJ|_v$. In what follows we will freely use the notation of [40], p. 82, which is explained there in details and drop the index "v", as it is clear that all objects are local ones at a non-split, archimedean place.
Let us begin with the non-split case, i.e., with $G' = G'_v$ and a representation $A_{\mathfrak{q}'_{\underline{k}}}(\lambda)$ index by $\underline{k} = [k_0, k_1, ..., k_r]$ as in Section 4.2. Recall the Levi subgroup $L' = L'_{\underline{k}} \cong GL_{k_0}(\mathbb{H}) \times \prod_{i=1}^{r} GL_{k_i}(\mathbb{C})$ defined in Section 4.4. It contains a maximally split θ'-stable Cartan subgroup T^+A^d. In fact, for the Lie algebras we may take

$$\mathfrak{t}^+ = Lie(T^+) = \mathfrak{t}' \qquad \text{and} \qquad \mathfrak{a}^d = Lie(A^d) = \mathfrak{a}',$$

where $\mathfrak{t}'$ and $\mathfrak{a}'$ are as in Section 4.2. As in [40], p. 82, we let M^dA^d be the Langlands decomposition of the centralizer of A^d in $G' = GL_k(\mathbb{H})$ and obtain

$$M^dA^d \cong \prod_{j=1}^{k} GL_1(\mathbb{H}),$$

so $M^d \cong \prod_{j=1}^{k} SL_1(\mathbb{H})$ and let P' be a parabolic subgroup containing M^dA^d as Levi factor. It is a minimal parabolic subgroup. The character ν^d defined on [40], p.82, is given as

$$\begin{aligned} \nu^d &= (\rho_{\mathfrak{gl}_{k_0}(\mathbb{H})}, \rho_{\mathfrak{gl}_{k_1}(\mathbb{C})}, ..., \rho_{\mathfrak{gl}_{k_r}(\mathbb{C})}) \\ &=: \nu'_{\underline{k}}, \end{aligned}$$

where $\rho_\bullet$ is the smallest algebraically integral element in the interior of the dominant Weyl chamber of $\bullet$. We still need to describe the representation σ^d of M^d, defined abstractly on p. 82 of [40]. Therefore, we recall three facts. Firstly, as we may assume that λ is self-dual, cf. Section 4.4, $\lambda|_{\mathfrak{t}^+} = \lambda = (\lambda_1, ..., \lambda_k)$ in the coordinates given by the functionals e_j, cf. Section 4.2. Secondly, ρ^+ being defined as the half sum of positive roots of $\mathfrak{t}^+ = \mathfrak{t}'$ in $\mathfrak{m}^d \cap \mathfrak{l}'$ equals, $\rho^+ = (0, ..., 0, 1, ..., 1)$, the "1's" starting from the $k - k_0 + 1$-st entry.

Thirdly, the highest weight of σ^d with respect to the system $\Delta(\mathfrak{m}^d_{\mathbb{C}}, \mathfrak{t}'_{\mathbb{C}})$ equals the Harish-Chandra parameter of σ^d subtracted by $\rho_{\mathfrak{m}^d} = (1, 1, ..., 1)$. Having collected this information, and writing $\rho(\mathfrak{u}') = (\rho_1^{\mathfrak{u}'}, ..., \rho_k^{\mathfrak{u}'})$ for the half-sum of the roots appearing in $\mathfrak{u}' = \mathfrak{u}'_{\underline{k}}$, counted with multiplicity, [40], (6.13) tells us that

$$\sigma^d \cong \bigotimes_{j=1}^{k-k_0} \mathrm{Sym}^{\lambda_j+\rho_j^{\mathfrak{u}'}-1}\mathbb{C}^2 \otimes \bigotimes_{j=1}^{k_0} \mathrm{Sym}^{\lambda_{j+k-k_0}+\rho^{\mathfrak{u}'}_{j+k-k_0}}\mathbb{C}^2.$$

However, by Lem. 4.1 $\rho^{\mathfrak{u}'}_{j+k-k_0} = 0$ for $1 \leq j \leq k_0$ and by (4.8) necessarily also $\lambda_{j+k-k_0} = 0$ for $1 \leq j \leq k_0$. So,

$$\begin{aligned} \sigma^d &\cong \bigotimes_{j=1}^{k-k_0} \mathrm{Sym}^{\lambda_j+\rho_j^{\mathfrak{u}'}-1}\mathbb{C}^2 \\ &= \bigotimes_{j=1}^{k-k_0} F(0, \lambda_j + \rho_j^{\mathfrak{u}'}) \\ &=: \sigma'_{\lambda,\underline{k}}, \end{aligned}$$

where $F(0, \lambda_j + \rho_j^{\mathfrak{u}'})$ is the notation used in the previous section. By [40], Thm. 6.16, $A_{\mathfrak{q}'_{\underline{k}}}(\lambda)$ is isomorphic to the unique irreducible quotient $J(P', \sigma'_{\lambda,\underline{k}}, \nu'_{\underline{k}})$ of the parabolically induced representation

$$\mathrm{Ind}_{P'}^{G'}[\sigma'_{\lambda,\underline{k}} \otimes \nu'_{\underline{k}}].$$

Next, we treat the split case, i.e., $G = GL_n(\mathbb{R})$, $n = 2k \geq 2$, and a representation $A_{\mathfrak{q}_{\underline{n}}}(\lambda)$ parameterized by a partition $\underline{n} = [n_0, n_1, ..., n_r]$ as in Section 4.3. From 4.4 we have a Levi subgroup $L = L_{\underline{n}} \cong GL_{n_0}(\mathbb{R}) \times \prod_{i=1}^r GL_{k_i}(\mathbb{C})$ containing a maximally split θ-stable Cartan subgroup which we again denote by T^+A^d, following [40], p.82. The Lie algebras of its factors satisfy

$$\mathfrak{t}^+ = Lie(T^+) \hookrightarrow \mathfrak{t} \qquad \text{and} \qquad \mathfrak{a} \hookrightarrow \mathfrak{a}^d = Lie(A^d),$$

where $\mathfrak{t}$ and $\mathfrak{a}$ are as in Section 4.3. The Langlands decomposition of the centralizer of A^d in $G = GL_n(\mathbb{R})$ is given by

$$M^d A^d \cong \prod_{j=1}^{k-k_0} GL_2(\mathbb{R}) \times \prod_{j=1}^{n_0} GL_1(\mathbb{R}),$$

so $M^d \cong \prod_{j=1}^{k-k_0} SL_2^{\pm}(\mathbb{R}) \times \prod_{j=1}^{n_0}\{\pm 1\}$ and we let P be a parabolic subgroup containing $M^d A^d$ as Levi factor. The character ν^d defined on [40], p. 82, takes the form

$$\begin{aligned} \nu^d &= (\rho_{\mathfrak{gl}_{n_0}(\mathbb{R})}, \rho_{\mathfrak{gl}_{k_1}(\mathbb{C})}, ..., \rho_{\mathfrak{gl}_{k_r}(\mathbb{C})}) \\ &=: \nu_{\underline{n}}. \end{aligned}$$

Next, we turn to the discrete series representation σ^d of M^d. According to [40], (6.13) the Harish-Chandra parameter of σ^d is

$$\rho^+ + \lambda|_{\mathfrak{t}^+} + \rho(\mathfrak{u}_{\underline{n}}).$$

Observing that $\mathfrak{m}^d \cap \mathfrak{l} \cong \mathfrak{t}^+$, implies $\rho^+ = (0, 0, ..., 0)$. Furthermore, $\lambda|_{\mathfrak{t}^+} = \lambda = (\lambda_1, ..., \lambda_{k-k_0}, 0, ...0)$, so we obtain that σ^d has Harish-Chandra parameter $\lambda|_{\mathfrak{t}^+} + \rho(\mathfrak{u}_{\underline{n}})$. The lowest $M^d \cap K$-type of σ^d, cf. [40] (6.14), affords that $m \in \prod_{j=1}^{n_0}\{\pm 1\}$ acts by $\mathrm{sgn}(\det(m))^{e(\lambda)}$ for some $e(\lambda) \in \{0, 1\}$. If we observe that the roots appearing in $\mathfrak{u}_{\underline{n}}$ and $\mathfrak{u}'_{\underline{k}}$ are the same, by Lemmas 4.1 and 4.5, this defines σ^d uniquely as

$$\begin{aligned} \sigma^d &\cong \bigotimes_{j=1}^{k-k_0} D(0, \lambda_j + \rho_j^{\mathfrak{u}'}) \otimes \mathrm{sgn}^{e(\lambda)} \\ &=: \sigma_{\lambda,\underline{n}} \end{aligned}$$

in the notation introduced in the previous section. If we use once more [40], Thm. 6.16, we obtain that $A_{\mathfrak{q}_{\underline{n}}}(\lambda)$ is isomorphic to the unique irreducible quotient $J(P,\sigma_{\lambda,\underline{n}},\nu_{\underline{n}})$ of the parabolically induced representation

$$\mathrm{Ind}_P^G[\sigma_{\lambda,\underline{n}}\otimes\nu_{\underline{n}}].$$

But now, as $|LJ|$ commutes with parabolic induction and forming tensor products, Lem. 5.1 together with induction by stages shows that

$$|LJ|(J(P,\sigma_{\lambda,\underline{n}},\nu_{\underline{n}}))=J(P',\sigma'_{\lambda,\underline{k}},\nu'_{\underline{k}}).$$

But this implies that

$$|LJ|(A_{\mathfrak{q}_{\underline{n}}}(\lambda)\otimes\mathrm{sgn}^{\varepsilon})=A_{\mathfrak{q}'_{\underline{k}}}(\lambda),$$

where $\underline{n}=[n_0,n_1,...,n_r]$ and $\underline{k}=[k_0,k_1,...,k_r]$ with $n_i=2k_i$, $0\leq i\leq r$. In particular, following the characterization of the cohomological, irreducible, unitary dual of G' and G in our Prop. 4.9 and 4.12, $|LJ|$ defines a surjection

$$\mathrm{Coh}_\mu(G)\xrightarrow{|LJ|}\mathrel{\mkern-14mu}\rightarrow\mathrm{Coh}_\mu(G')$$

for all highest weight representations $E=E_\mu$. Furthermore, by Thm. 4.9 (4) and 4.12 (3), $|LJ|$ maps tempered cohomological representations to tempered cohomological representations. This shows the theorem. □

Remark 5.3. The fibers of $|LJ|_v$ over a representation $A_{\mathfrak{q}'}(\lambda)$ can be explicitly described using (4.4).

5.4. **An illustrative example.** Let us exemplify Thm. 5.2 by letting $k=2$, so $n=4$, and taking $E=\mathbb{C}$ to be the trivial representation, i.e., $\lambda=0$. We obtain the following two tables, where Q is the standard parabolic subgroup of $G=GL_4(\mathbb{R})$ with Levi factor $GL_2(\mathbb{R})\times GL_2(\mathbb{R})$ and P' is as in the proof of Thm. 5.2.

$\underline{n}$	$A_{\mathfrak{q}_{\underline{n}}}(0)$	$A_{\mathfrak{q}_{\underline{n}}}(0)\otimes\mathrm{sgn}$
$[0,2,2]$	$\mathrm{Ind}_Q^G[D(0,3)\otimes D(0,1)]$	$\mathrm{Ind}_Q^G[D(0,3)\otimes D(0,1)]$
$[2,2]$	$\mathrm{Ind}_Q^G[D(0,3)\otimes \mathbf{1}_{GL_2(\mathbb{R})}]$	$\mathrm{Ind}_Q^G[D(0,3)\otimes \mathrm{sgn}]$
$[0,4]$	$J(Q,D(-1,2)\otimes D(1,2))$	$J(Q,D(-1,2)\otimes D(1,2))$
$[4]$	$\mathbf{1}_{GL_4(\mathbb{R})}$	sgn

TABLE 1. $\mathrm{Coh}_0(GL_4(\mathbb{R}))$

$\underline{k}$	$A_{\mathfrak{q}'_{\underline{k}}}(0)$
$[0,1,1]$	$\mathrm{Ind}_{P'}^{G'}[F(0,3)\otimes F(0,1)]$
$[1,1]$	$\mathrm{Ind}_{P'}^{G'}[F(0,3)\otimes \mathbf{1}_{GL_1(\mathbb{H})}]$
$[0,2]$	$J(P',F(-1,2)\otimes F(1,2))$
$[2]$	$\mathbf{1}_{GL_2(\mathbb{H})}$

TABLE 2. $\mathrm{Coh}_0(GL_2(\mathbb{H}))$

Since $F(0,1)=\mathbf{1}_{GL_1(\mathbb{H})}$, the representations in the first two lines of Table 2 coincide as predicted. Furthermore, the tempered representations are given by the first rows of Table 1 and 2, i.e., by the partitions $[0,2,2]$ and $[0,1,1]$. Applying Lem. 5.1 directly to Table 1 and noting that $\mathrm{Ind}_Q^G[D(0,3)\otimes\mathbf{1}_{GL_2(\mathbb{R})}]$ is isomorphic to the irreducible quotient of $\mathrm{Ind}_P^G[D(0,3)\otimes\mathbf{1}_{GL_1(\mathbb{R})}|\cdot|^{1/2}\otimes\mathbf{1}_{GL_1(\mathbb{R})}|\cdot|^{-1/2}]$, where P is as in the proof of Thm. 5.2, reproves the other assertions in this example.

5.5. **The essentially-tempered representations.** There is a particularly important subclass of irreducible, cohomological representations, namely those which are *essentially tempered*. Therefore recall that an irreducible admissible representation Π'_v of G'_v, $v \in V_\infty$, is called *essentially tempered*, if it is of the form $\Pi'_v \cong \tilde{\Pi}'_v \otimes |\det'|^{s_v}$, with $\tilde{\Pi}'_v$ being an irreducible, unitary, tempered representation and $s_v \in \mathbb{C}$. A representation $\Pi'_\infty = \bigotimes_{v\in V_\infty} \Pi'_v$ of G'_∞ is analogously called essentially tempered if all its local factors Π'_v are. The analog definition applies to representations of G_v when removing the prime.

According to Thm. 5.2 and the definition of $|LJ|_v$, a cohomological, irreducible, admissible, essentially tempered representation Π_v of G_v is mapped via $|LJ|_v$ onto a cohomological, irreducible, admissible, essentially tempered representation Π'_v of G'_v. Let now $v \in V_\infty$ be again non-split and let us shortly describe these representations a little bit more explicitly.

So, let $E = E_{\mu_v}$ be a highest weight representation as in Section 2.3. For sake of simplicity, we will again drop the subscript "v" in this subsection for all local objects. By the same reason as in Section 4.4 we see that, in order to admit a cohomological, essentially tempered, irreducible admissible representation Π' of G', it is necessary for the coefficient system E_μ to be essentially self-dual. We may hence assume without loss of generality that from now on in this subsection E_μ is essentially self-dual writing $E_\mu = E_\mu^\vee \otimes \det^w$, $w \in \mathbb{Z}$. Otherwise put, if $\mu = (\mu_1, ..., \mu_n)$ is the highest weight of E_μ,

$$\mu_i + \mu_{n-i+1} = w \quad \text{for} \quad 1 \le i \le n.$$

Let $\ell := (\ell_1, ..., \ell_k)$ with

$$\ell_i := \mu_i - \mu_{n-i+1} + (n - 2i + 1) = w - 2\mu_{n-i+1} + (n - 2i + 1).$$

We obtain that $\ell_1 > \ell_2 > ... > \ell_k \ge 1$. By Thm. 4.9 (4) and Borel–Wallach [7], I, Thm. 5.3, the unique, irreducible, admissible, essentially tempered representation of G', which has non-trivial $(\mathfrak{g}', K'^\circ)$-cohomology twisted by E_μ is

$$A_{\mathfrak{q}'_{\underline{k}}}(\lambda) \otimes \det'^{-w/2},$$

where $\underline{k} = [0, 1, 1, ..., 1]$. It is furthermore a consequence of the proof of Thm. 5.2, that this latter representation is isomorphic to the unique irreducible quotient of

$$\mathrm{Ind}_{P'}^{G'}[F(w, \ell_1) \otimes ... \otimes F(w, \ell_k)].$$

Now, [17], Thm. 1, p. 198, implies that this normalized induced representation is already irreducible itself, whence

$$J'(w, \ell) := A_{\mathfrak{q}'_{\underline{k}}}(\lambda) \otimes \det'^{-w/2} \cong \mathrm{Ind}_{P'}^{G'}[F(w, \ell_1) \otimes ... \otimes F(w, \ell_k)]$$

is the unique, irreducible, admissible, essentially tempered representation of G', which has non-trivial $(\mathfrak{g}', K'^\circ)$-cohomology twisted by E_μ. Furthermore,

$$H^q(\mathfrak{gl}_k(\mathbb{H}), Sp(k)\mathbb{R}_+, J'(w, \ell) \otimes E_\mu) \cong \bigwedge^{q-k(k-1)} \mathbb{C}^{k-1},$$

which follows from Thm. 4.9 (3) or directly by [7] III. Thm 5.1.

The split case $G = G_v = GL_n(\mathbb{R})$, $n = 2k$, is treated in analogy. By Thm. 4.12 and [7], I, Thm. 5.3, the unique, irreducible, admissible, essentially tempered representation of G, which has non-trivial $(\mathfrak{g}, K^\circ)$-cohomology twisted by E_μ is

$$A_{\mathfrak{q}_{\underline{n}}}(\lambda) \otimes |\det|^{-w/2} \cong A_{\mathfrak{q}_{\underline{n}}}(\lambda) \otimes \mathrm{sgn}\, |\det|^{-w/2},$$

where $\underline{n} = [0, 2, 2, ..., 2]$. As shown in the proof of Thm. 5.2, this latter representation is isomorphic to the unique irreducible quotient of

$$\mathrm{Ind}_P^G[D(w, \ell_1) \otimes ... \otimes D(w, \ell_k)],$$

which turns out to be irreducible itself, too, cf. [17], Thm. 1, p. 198. Hence,

$$J(w, \ell) := A_{\mathfrak{q}_{\underline{n}}}(\lambda) \otimes |\det|^{-w/2} \cong \mathrm{Ind}_P^G[D(w, \ell_1) \otimes ... \otimes D(w, \ell_k)]$$

is the unique, irreducible, admissible, essentially tempered representation of G, which has non-trivial $(\mathfrak{g}, K^\circ)$-cohomology twisted by E_μ. We compute,

$$H^q(\mathfrak{gl}_n(\mathbb{R}), SO(n)\mathbb{R}_+, J(w,\ell) \otimes E_\mu) \cong \mathbb{C}^2 \otimes \bigwedge^{q-k^2} \mathbb{C}^{k-1}$$

which follows again from [7] III. Thm 5.1 or directly using [7], III. Thm. 3.3. Observe that there is a factor $\mathbb{C}^2$ appearing in the formula. This is due to the fact that we calculated $(\mathfrak{gl}_n(\mathbb{R}), SO(n)\mathbb{R}_+)$-cohomology (as in Mahnkopf [25], 3.1.2), and not $(\mathfrak{gl}_n(\mathbb{R}), O(n)\mathbb{R}_+)$-cohomology (as it was done by Clozel in [8], Lemme 3.14). The reason for that will become clear after Thm. 6.3. Compare this also to [25], p. 590.

5.6. **The global Jacquet-Langlands map.** There is also a global version of the Jacquet-Langlands map, developed by Badulescu in [1], sect. 5, and by Badulescu–Renard in [2], sect. 18. (For $m = 1$ see also Harris–Taylor [18], VI.1) We denote it here by JL (because it goes in the direction different to the one of $|LJ|_v$). It is uniquely determined as being the map satisfying the conditions of the following theorem:

Theorem 5.4 ([2], Thm. 18.1 & Prop. 18.2)**.** *The map JL is the unique injection from the set of all unitary discrete series representations $\tilde{\Pi}'$ of $G'(\mathbb{A})$ into the set of all unitary discrete series representations $\tilde{\Pi}$ of $G(\mathbb{A})$ such that*

$$|LJ|_v(JL(\tilde{\Pi}')_v) = \tilde{\Pi}'_v$$

at all places $v \in V$.
Moreover, if $JL(\tilde{\Pi}')$ is unitary cuspidal, then so is $\tilde{\Pi}'$.

We remark that the condition $|LJ|_v(JL(\tilde{\Pi}')_v) = \tilde{\Pi}'_v$ implies that in particular $JL(\tilde{\Pi}')_v = \tilde{\Pi}'_v$ at all split places v. In accordance with our local definition (cf. Section 5.1), we define for $\Pi' = \tilde{\Pi}' \otimes |\det'|^s \in \mathscr{D}(G')$:

$$JL(\Pi') := JL(\tilde{\Pi}') \otimes |\det|^s.$$

Corollary 5.5. *The so extended global Jacquet-Langlands map satisfies $JL(\Pi') \in \mathscr{D}(G)$. Furthermore, all assertions of Thm. 5.4 hold unchanged (omitting the word "unitary").*

6. Algebraic and regular algebraic representations of $G'(\mathbb{A})$

6.1. **The Local Langlands Classification.** Let $W_{\mathbb{K}}$ be the local Weil group of $\mathbb{K} = \mathbb{R}, \mathbb{C}$, cf. Tate [37], 1.4.3. It can be defined as follows:

$$W_{\mathbb{K}} = \begin{cases} \mathbb{C}^* & \text{if } \mathbb{K} = \mathbb{C} \\ \mathbb{C}^* \cup j\mathbb{C}^* & \text{if } \mathbb{K} = \mathbb{R}, \end{cases}$$

where in the second case $j^2 = -1$ and $jzj^{-1} = \overline{z}$, for all $z \in \mathbb{C}^*$.

It was proved by Langlands in [22] that there is a canonical bijection between the class of irreducible admissible representations π of $GL_n(\mathbb{K})$ and the class of semi-simple, complex, n-dimensional representations τ of $W_{\mathbb{K}}$, $\mathbb{K} = \mathbb{R}, \mathbb{C}$. We denote this correspondence by

$$\pi \leftrightarrow \tau = \tau(\pi)$$

and call $\tau(\pi)$ the *Langlands parameter* of π. If $\mathbb{K} = \mathbb{R}$ we can furthermore restrict $\tau(\pi)$ to the connected component of the local Weil group $W_{\mathbb{R}}^\circ \cong \mathbb{C}^*$, which gives us an n-dimensional, complex representation of $\mathbb{C}^*$, again denoted by $\tau(\pi)$. This enables us to view $\tau(\pi)$ – no matter if $\mathbb{K} = \mathbb{R}$ or $\mathbb{K} = \mathbb{C}$ – as a representation

$$\tau(\pi) : \mathbb{C}^* \to GL_n(\mathbb{C}).$$

If $\mathbb{K} = \mathbb{R}$ and $\pi \cong D(l) \otimes |\det|^s$ (notation as in Section 5.2) for some integer $l \geq 1$ and $s \in \mathbb{C}$, then the restriction of $\tau(\pi)$ to $\mathbb{C}^*$ is explicitly given by

$$z \mapsto (z^{\frac{l}{2}+s}(\overline{z})^{-\frac{l}{2}+s}) \oplus (z^{-\frac{l}{2}+s}(\overline{z})^{\frac{l}{2}+s}).$$

6.2. **The split case.** We will now recall the definition of an algebraic and a regular algebraic automorphic representation of GL_n/F, following Clozel.

Definition 6.1 ([8], Def. 1.8 & Def. 3.12)**.** Let $\Pi \in \mathscr{D}(G)$ with archimedean component $\Pi_\infty = \bigotimes_{v\in V_\infty} \Pi_v$.

(1) The representation Π is called *algebraic*, if at all places $v \in V_\infty$

$$\Pi_v \leftrightarrow \tau(\Pi_v) = \chi_{1,v} \oplus ... \oplus \chi_{n,v},$$

where $\chi_{i,v}|\cdot|_{\mathbb{C}}^{\frac{1-n}{2}}$, $1 \leq i \leq n$, is an algebraic character and $|\cdot|_{\mathbb{C}}$ denotes the usual normalized absolute value on $\mathbb{C}$.

(2) An algebraic representation Π is furthermore called *regular algebraic*, if the infinitesimal character χ_{Π_∞} of Π_∞ is regular, i.e., inside the positive, open Weyl chamber.

Remark 6.2. A character $\chi_{i,v}|\cdot|_{\mathbb{C}}^{\frac{1-n}{2}}$ being algebraic means that

$$\chi_{i,v}(z) = z^{p_{i,v}}(\overline{z})^{q_{i,v}}$$

with $p_{i,v}, q_{i,v} \in \frac{n-1}{2} + \mathbb{Z}$ for all $v \in V_\infty$ and $1 \leq i \leq n$. If v is a real place, we can even more suppose that $\tau(\Pi_v)$ is given by

$$z \mapsto (z^{p_{1,v}}(\overline{z})^{p_{2,v}}) \oplus (z^{p_{2,v}}(\overline{z})^{p_{1,v}}) \oplus ... \oplus ((z\overline{z})^{p_{2r+1,v}}) \oplus ... \oplus ((z\overline{z})^{p_{n,v}}).$$

We let $p_v = (p_{1,v}, ..., p_{n,v}) \in (\frac{n-1}{2} + \mathbb{Z})^n$ for all $v \in V_\infty$. The definition of Π being regular algebraic can now be equivalently reformulated by saying that for all $v \in V_\infty$, $p_{i,v} \neq p_{j,v}$, $1 \leq i < j \leq n$. (This is actually the original definition of Clozel, cf. [8], Def. 3.12.)

There is also the following, very useful, equivalent description of regular algebraic, cuspidal representations Π, which is a consequence of Clozel [8], Lemme 3.14.

Theorem 6.3. *Let $\Pi \in \mathscr{D}(G)$ and assume Π is cuspidal. Then the following are equivalent:*

(i) Π *regular algebraic.*

(ii) Π_∞ *is cohomological and essentially tempered.*

Sketch of proof. By Clozel [8], Lemme 3.14, a cuspidal automorphic representation Π is regular algebraic if and only if Π_∞ is cohomological. So, we only need to show that the archimedean component Π_∞ of a cuspidal *and* cohomological representation Π is automatically essentially tempered. This is actually well-known, so we sketch the argument here. For $n = 1$, this is obvious. If $n \geq 2$, then by Shalika [34], corollary on p. 190, Π_v is a generic representation of G_v for all $v \in V$. But a generic, cohomological, irreducible admissible representation of G_v, $v \in V_\infty$, is essentially tempered. This shows the claim. □

Remark 6.4. We want to point out that it is due to the fact that we calculate $(\mathfrak{gl}_n(\mathbb{R}), SO(n)\mathbb{R}_+)$ - cohomology at a split real place v that we can formulate the theorem in that way. If we calculated $(\mathfrak{gl}_n(\mathbb{R}), O(n)\mathbb{R}_+)$-cohomology instead, if n is odd, one might have to twist Π_v with the sign-character in order to get non-vanishing cohomology, cf. [8], Lem. 3.14.

Moreover, we remark that the cuspidality assumption on Π cannot be removed in Thm. 6.3. To see this, consider the following counter-example:

Example 6.5. Let $F = \mathbb{Q}$ and take π to be a cuspidal automorphic representation of $GL_2(\mathbb{A})$ with $\pi_\infty \cong D(2)$ (notation as in Section 5.2). If we put moreover $n = 4$, then the global induced representation

$$\mathrm{Ind}_{GL_2(\mathbb{A})\times GL_2(\mathbb{A})}^{GL_4(\mathbb{A})}[\pi|\det|^{1/2} \otimes \pi|\det|^{-1/2}]$$

has a unique irreducible quotient Π, which is a residual representation in $\mathscr{D}(GL_4)$, see Mœglin–Waldspurger [26]. At infinity, Π_∞ is the Langlands quotient of

$$\mathrm{Ind}_{GL_2(\mathbb{R})\times GL_2(\mathbb{R})}^{GL_4(\mathbb{R})}[D(2)|\det|^{1/2} \otimes D(2)|\det|^{-1/2}].$$

Therefore, by Section 6.1, $\tau(\Pi_\infty)$ is given by the character

$$z \mapsto (z^{\frac{3}{2}}(\overline{z})^{-\frac{1}{2}}) \oplus (z^{-\frac{1}{2}}(\overline{z})^{\frac{3}{2}}) \oplus (z^{\frac{1}{2}}(\overline{z})^{-\frac{3}{2}}) \oplus (z^{-\frac{3}{2}}(\overline{z})^{\frac{1}{2}}),$$

whose exponents are all in $\frac{1}{2}+\mathbb{Z}$ and pairwise different, whence Π is regular algebraic by Rem. 6.2. Its archimedean component Π_∞ is even cohomological with respect to the trivial representation: One may check using Thm. 5.2, resp. Table 1, that

$$H^q(\mathfrak{gl}_4(\mathbb{R}), SO(4)\mathbb{R}^+, \Pi_\infty \otimes \mathbb{C}) \neq 0$$

in degrees $q = 3, 6$. But clearly, Π_∞ is not tempered. Indeed, $\Pi_\infty = A_{\mathfrak{q}_{\underline{n}}}(\lambda)$ with $\lambda = 0$ and $\underline{n} = [0, 4]$, which is not tempered by Thm. 4.12 (3).

Remark 6.6. If $n = 2k$ and $v \in V_\infty$ is a real place, the local component Π_v of a regular algebraic cuspidal representation $\Pi \in \mathscr{D}(G)$ is necessarily of the form $\Pi_v \cong J(w_v, \ell_v)$, for some integer w_v and tuple ℓ_v as in Section 5.5. In view of Thm. 6.3 this follows from Thm. 4.12 (resp., Section 5.5).

Before we turn to the case of G', let us also recall Clozel's "Lemme de pureté", [8], Lemme 4.9:

Lemma 6.7. *If $\Pi \in \mathscr{D}(G)$ is algebraic and cuspidal, then there is a $\mathsf{w} \in \mathbb{Z}$ such that for all $v \in V_\infty$, the algebraic characters of $\mathbb{C}^*$ associated to $\Pi_v|\cdot|_v^{\frac{1-n}{2}}$ are of the form $z \mapsto z^p(\overline{z})^q$ with $p + q = \mathsf{w}$.*

6.3. **The general case.** Motivated by the above definition for the split case (i.e., $D = F$) we extend it to the more general case of G'.

Definition 6.8. Let $\Pi' \in \mathscr{D}(G')$.

(1) The representation Π' is called *algebraic*, if $JL(\Pi') \in \mathscr{D}(G)$ is algebraic.
(2) An algebraic representation Π' is called *regular algebraic*, if $JL(\Pi') \in \mathscr{D}(G)$ is regular algebraic.

We will now see that our definition goes very well with Thm. 6.3.

Theorem 6.9. *Let $\Pi' \in \mathscr{D}(G')$ and assume that $JL(\Pi')$ is cuspidal. Then the following are equivalent:*

(i) *Π' is regular algebraic.*
(ii) *Π'_∞ is cohomological and essentially tempered.*

Proof. (i)⇒(ii): By assumption $JL(\Pi')$ is a cuspidal, regular algebraic representation of $G(\mathbb{A})$. So, Thm. 6.3 implies that $JL(\Pi')_\infty = \bigotimes_{v \in V_\infty} JL(\Pi')_v$ is cohomological and essentially tempered. By the description of the global Jacquet-Langlands map, cf. Thm. 5.4 and Cor. 5.5, there is an isomorphism $JL(\Pi')_v \cong \Pi'_v$ at all split places $v \in V$. In particular, Π'_v is proved to be cohomological and essentially tempered at all split places $v \in V_\infty$. So, let us assume from now on that v is a real, non-split place. Again by Thm. 5.4 and Cor. 5.5 we obtain

$$\Pi'_v \cong |LJ|_v(JL(\Pi')_v).$$

As v is supposed to be non-split, we have $n = 2k$ and so Thm. 6.3 (see also Rem. 6.6) implies that

$$JL(\Pi')_v \cong J(w_v, \ell_v),$$

for some integer w_v and k-tuple ℓ_v as in Section 5.5. But now Thm. 5.2 shows that $|LJ|_v(J(w_v, \ell_v)) \cong J'(w_v, \ell_v)$. Putting the pieces together we finally see that

$$\Pi'_v \cong J'(w_v, \ell_v).$$

According to Thm. 4.9 (resp., Section 5.5), Π'_v is therefore cohomological and essentially tempered and hence so is $\Pi'_\infty = \bigotimes_{v \in V_\infty} \Pi'_v$.

(ii)⇒(i): Assume that Π'_∞ is cohomological and essentially tempered. By definition, cf. Def. 6.8, we need to show that $JL(\Pi')$ is regular algebraic. As this is a local condition at the archimedean places, Thm. 6.3 together with the standing assumption that $JL(\Pi')$ is cuspidal, permits us again to focus on the non-split archimedean places $v \in V_\infty$. Let $v \in V_\infty$ be such a place and let E_{μ_v}, $\mu_v = (\mu_{v,1}, ..., \mu_{v,n})$, be a highest weight representation of $G'_v = GL_k(\mathbb{H})$ with respect to which Π'_v is cohomological. We know from Section 5.5 that E_{μ_v} is necessarily essentially self-dual, so there is an integer $w_v \in \mathbb{Z}$ such that $w_v = \mu_{v,i} + \mu_{v,n-i+1}$, $1 \leq i \leq n$. As Π'_v is furthermore assumed to be essentially tempered, Thm. 4.9, resp., our explanations of Section 5.5, imply that

$$\Pi'_v \cong J'(w_v, \ell_v),$$

the k-tuple $\ell_v = (\ell_{1,v}, ..., \ell_{k,v})$ being defined as in Section 5.5. By the description of the global Jacquet-Langlands map, we hence obtain

$$J'(w_v, \ell_v) \cong |LJ|_v(JL(\Pi')_v).$$

Next, recall that $JL(\Pi')_v$ – being the local component of a cuspidal representation – must be generic (see Shalika [34], corollary on p. 190). Therefore $JL(\Pi')_v$ is induced from representations in the (limits of) essentially discrete series, cf. Vogan [38], Thm. 6.2. But since $|LJ|_v$ commutes with induction and tensor products, these representations must be all in $U_{cp}(GL_2(\mathbb{R}))$ and so Lem. 5.1 shows that $JL(\Pi')_v$ is induced from the representations $D(w_v, \ell_{i,v})$, $1 \leq i \leq k$. Whence we have by construction

$$JL(\Pi')_v \cong J(w_v, \ell_v)$$

and so Thm. 4.12 (resp., Section 5.5) yields that

$$JL(\Pi')_\infty = \bigotimes_{v \in V_\infty} JL(\Pi')_v$$

is cohomological and essentially tempered. By Thm. 6.3, $JL(\Pi')$ is therefore regular algebraic. Now the proof is complete. $\square$

Remark 6.10. Implication (ii)⇒(i) does not seem to need the assumption that $JL(\Pi')$ is cuspidal. First, observe that – assuming the validity of (ii) – Π' is necessarily cuspidal by Wallach [42], Thm. 4.3. See also Clozel [9], Prop. 4.10. By the same references, $JL(\Pi')$ is cuspidal, if there is a split archimedean place $v \in V_\infty$. It therefore suffices to prove the implication (ii)⇒(i) under the assumption that there are only non-split places $v \in V_\infty$. It seems that Π'_v, being cohomological and essentially tempered, satisfies

$$JL(\Pi')_v \cong A_{\mathfrak{q}_{\underline{n}}}(\lambda) \otimes \mathrm{sgn}^\varepsilon |\mathrm{det}|^{-w/2}, \qquad (\heartsuit)$$

with $\underline{n} = [0, 2, 2, ..., 2]$ or $\underline{n} = [2, 2, ..., 2]$. If $\underline{n} = [0, 2, 2, ..., 2]$, then $JL(\Pi')_v$ is essentially tempered and we are again in the situation considered in Thm. 6.9. If $\underline{n} = [2, 2, ..., 2]$, then the local Langlands parameter of $JL(\Pi')_v$ is given by

$$(z^{\frac{\ell_1 - w}{2}} (\overline{z})^{\frac{-\ell_1 - w}{2}}) \oplus (z^{\frac{-\ell_1 - w}{2}} (\overline{z})^{\frac{\ell_1 - w}{2}}) \oplus ... \oplus (z^{\frac{\ell_{k-1} - w}{2}} (\overline{z})^{\frac{-\ell_{k-1} - w}{2}}) \oplus (z^{\frac{-\ell_{k-1} - w}{2}} (\overline{z})^{\frac{\ell_{k-1} - w}{2}}) \oplus (z\overline{z})^{\frac{\ell_k - w}{2}} \oplus (z\overline{z})^{\frac{-\ell_k - w}{2}},$$

with $\ell = (\ell_1, ..., \ell_k)$ and w as in Section 5.5; in particular, $\ell_k = 1$. Indeed, for $A_{\mathfrak{q}_{\underline{n}}}(\lambda)$ with $\underline{n} = [2, 2, ..., 2]$ to exist, we must have $\lambda_k = 0$ by (4.8). As furthermore $\rho_k^{\mathfrak{u}_{\underline{n}}} = 0$, the representation giving $(z\overline{z})^{\frac{\ell_k - w}{2}} \oplus (z\overline{z})^{\frac{-\ell_k - w}{2}}$ is $|\det|^{-w/2}$, which shows $\ell_k = 1$. By the definition of the numbers ℓ_i we see that $\ell_i - w = 2(-\mu_{n-i+1} + k - i) + 1$ is odd, so $JL(\Pi')_v$ would be algebraic. But since also $\ell_1 > \ell_2 > ... > \ell_{k-1} > \ell_k = 1$, $JL(\Pi')_v$ would also be regular algebraic by Rem. 6.2. As a consequence, in order to show the implication (ii)⇒(i) without the assumption that $JL(\Pi')$ is cuspidal, it suffices to prove (♡).

In contrast, the implication (i)⇒(ii) in Thm. 6.9 fails, if one drops the assumption of the cuspidality of $JL(\Pi')$ – even if one supposes that Π' is cuspidal. This is shown in the next example:

Example 6.11 (Grobner [14], Thm. 4.1)**.** Let $F = \mathbb{Q}$, $m = 2$ and $d = 2$, i.e., G' is the group of $GL_2(D)$ over a quaternion division algebra D over $\mathbb{Q}$. Assume furthermore that the infinite place is non-split, so $G'_\infty = GL_2(\mathbb{H})$. Using the global Jacquet-Langlands map of [2], it was shown by the first named author in [14], Thm. 4.1, that for any integer $k \geq 0$, there is a cuspidal automorphic representation Π' of $G'(\mathbb{A})$ whose component at infinity Π'_∞ is given by the Langlands quotient of

$$\mathrm{Ind}_{GL_1(\mathbb{H}) \times GL_1(\mathbb{H})}^{GL_2(\mathbb{H})} [\mathrm{Sym}^{k+1}\mathbb{C}^2 \mathrm{det}'^{1/2} \otimes \mathrm{Sym}^{k+1}\mathbb{C}^2 \mathrm{det}'^{-1/2}].$$

Therefore, Π'_∞ is not essentially tempered, cf. [14], Prop. 3.5. (Although, if k is even, Π'_∞ is cohomological with respect to the highest weight representation E_μ of G'_∞ with $\mu = (\frac{k}{2}, \frac{k}{2}, -\frac{k}{2}, -\frac{k}{2})$; see again [14], Prop. 3.5, resp., our Thm. 4.9.)

But if k is chosen to be even, we claim that Π' is regular algebraic. In fact, the Langlands parameter $\tau(JL(\Pi')_\infty)$ is given by Section 6.1 as

$$z \mapsto (z^{\frac{(k+3)}{2}}(\overline{z})^{\frac{-(k+1)}{2}}) \oplus (z^{\frac{-(k+1)}{2}}(\overline{z})^{\frac{(k+3)}{2}}) \oplus (z^{\frac{(k+1)}{2}}(\overline{z})^{\frac{-(k+3)}{2}}) \oplus (z^{\frac{-(k+3)}{2}}(\overline{z})^{\frac{(k+1)}{2}}),$$

whose exponents are in $\frac{1}{2} + \mathbb{Z}$, if k is even. So, Π' is algebraic for even k. Furthermore, the numbers

$$p_{1,\infty} = \frac{(k+3)}{2}, p_{2,\infty} = \frac{-(k+1)}{2}, p_{3,\infty} = \frac{(k+1)}{2}, p_{4,\infty} = \frac{-(k+3)}{2}$$

can never be pairwise equal. Hence Π' is regular algebraic for even k by Rem. 6.2.

6.4. **A purity lemma.** We conclude this section by stating a generalization of Clozel's "Lemme de pureté".

Lemma 6.12. *If $\Pi' \in \mathscr{D}(G')$ is algebraic and $JL(\Pi')$ cuspidal, then there is a $\mathsf{w} \in \mathbb{Z}$ such that for all $v \in V_\infty$, the characters of $\mathbb{C}^*$ associated to $JL(\Pi')_v|\cdot|_v^{\frac{1-n}{2}}$ are of the form $z \mapsto z^p(\overline{z})^q$ with $p + q = \mathsf{w}$.*

Proof. This is an immediate consequence of the definition of Π' being algebraic and Lem. 6.7. ☐

Again, the cuspidality of $JL(\Pi')$ cannot be weakened to assuming that Π' is cuspidal, as the next example shows.

Example 6.13. Once more consider the representation Π' constructed in Example 6.11. It is cuspidal and algebraic, if k is even. Recalling also its Langlands parameter from Example 6.11, one checks that there is no integer $\mathsf{w} \in \mathbb{Z}$ representing all the sums of the exponents showing up in $\tau(JL(\Pi')_\infty)$.

7. SPACES OF AUTOMORPHIC COHOMOLOGY AND RATIONAL STRUCTURES

7.1. **Definition of the σ-twist, the rationality field and $\mathbb{F}$-structures.** Let ν be any representation of either $G'(\mathbb{A}_f)$ or G'_v on a complex vector space W. For $\sigma \in \mathrm{Aut}(\mathbb{C})$, we define the *$\sigma$-twist* ${}^\sigma\nu$ following Waldspurger [41], I.1: If W' is a $\mathbb{C}$-vector space which admits a σ-linear isomorphism $t : W \to W'$ then we set

$${}^\sigma\nu := t \circ \nu \circ t^{-1}.$$

This definition is independent of t and W' up to equivalence of representations. Further, if $\nu = \bigotimes_{v \in V_\infty} \nu_v$ is a representation of G'_∞, we let

$${}^\sigma\nu := \bigotimes_{v \in V_\infty} \nu_{\sigma^{-1}v},$$

interpreting $v \in V_\infty$ as an embedding of fields $v : F \hookrightarrow \mathbb{C}$.

Recall also the definition of the rationality field of a representation from [41], I.1. If ν is any of the representations considered above, then let $\mathfrak{S}(\nu)$ be the group of all automorphisms $\sigma \in \mathrm{Aut}(\mathbb{C})$ such that ${}^\sigma\nu \cong \nu$:

$$\mathfrak{S}(\nu) := \{\sigma \in \mathrm{Aut}(\mathbb{C}) | {}^\sigma\nu \cong \nu\}.$$

Then the *rationality field* $\mathbb{Q}(\nu)$ is defined as

$$\mathbb{Q}(\nu) := \{z \in \mathbb{C} | \sigma(z) = z \text{ for all } \sigma \in \mathfrak{S}(\nu)\}.$$

As a third ingredient we recall that a group representation ν on a $\mathbb{C}$-vector space W is said to be *defined over a subfield* $\mathbb{F} \subset \mathbb{C}$, if there is a $\mathbb{F}$-vector subspace $W_\mathbb{F} \subset W$, stable under the group action, and such that the canonical map $W_\mathbb{F} \otimes_\mathbb{F} \mathbb{C} \to W$ is an isomorphism. In this case, we also say that (ν, W) has an *$\mathbb{F}$-structure.*

In particular, if $E_\mu = \bigotimes_{v \in V_\infty} E_{\mu_v}$ is a highest weight representation of G'_∞, then the above definitions can be applied to E_μ. In this special case we will abbreviate

$$\mathbb{Q}(\mu) := \mathbb{Q}(E_\mu).$$

We obtain

Lemma 7.1 (Clozel [8], p.122)**.** *Let E_μ be any irreducible highest weight representation as in Section 2.3. Then $\mathbb{Q}(\mu)$ is a number field and E_μ is defined over its rationality field $\mathbb{Q}(\mu)$.*

Extending our notation, we let $\mathbb{Q}(\Pi')$ be the compositum of the fields $\mathbb{Q}(\mu)$ and $\mathbb{Q}(\Pi'_f)$,

$$\mathbb{Q}(\Pi') := \mathbb{Q}(\mu)\mathbb{Q}(\Pi'_f),$$

if Π' is a global automorphic representation of $G'(\mathbb{A})$, which has non-trivial cohomology with respect to E_μ.

7.2. **Rational structures on the cohomology of geometric spaces.** Consider now the quotient

$$S_{G'} := G'(F)\backslash G'(\mathbb{A})/K'^{\circ}_{\infty},$$

where we recall that the group $\mathbb{R}_+$ is "hidden" as a diagonal subgroup of K'°_{∞}. This space is the projective limit of finite disjoint unions of orbifolds, cf. Rohlfs [31], Prop. 1.9 together with Borel [4], Thm. 5.1. Let E_μ be a highest weight representation of G'_∞. It defines a sheaf $\mathcal{E}_\mu$ on $S_{G'}$, by letting $\mathcal{E}_\mu$ be the sheaf with espace étalé $G'(\mathbb{A})/K'^{\circ}_{\infty} \times_{G'(F)} E_\mu$ with the discrete topology on E_μ. Hence, the sheaf cohomology spaces

$$H^q(S_{G'}, \mathcal{E}_\mu)$$

are defined. These are $G'(\mathbb{A}_f)$-modules. Moreover, the sheaf-cohomology with compact support $H^q_c(S_{G'}, \mathcal{E}_\mu)$ is well-defined, too, and there is the natural map

$$H^q_c(S_{G'}, \mathcal{E}_\mu) \to H^q(S_{G'}, \mathcal{E}_\mu).$$

We denote its image, following Harder, by $H^q_!(S_{G'}, \mathcal{E}_\mu)$ and call it *interior cohomology.*

It will be important for us to interpret interior cohomology as the kernel of another map. Therefore, recall the adelic Borel-Serre Compactification $\overline{S}_{G'}$ of $S_{G'}$, and its basic properties. (For this we refer to Borel–Serre [6] as the original source and to Rohlfs [31] for the adelic setting.) It is a compact space with boundary $\partial(\overline{S}_{G'})$ and the inclusion $S_{G'} \hookrightarrow \overline{S}_{G'}$ is a homotopy equivalence. Furthermore, the sheaf $\mathcal{E}_\mu$ extends to a sheaf on $\overline{S}_{G'}$ and there is the natural restriction morphism of $G'(\mathbb{A}_f)$-modules

$$res^q : H^q(S_{G'}, \mathcal{E}_\mu) \cong H^q(\overline{S}_{G'}, \mathcal{E}_\mu) \to H^q(\partial(\overline{S}_{G'}), \mathcal{E}_\mu).$$

It is a basic fact that

$$H^q_!(S_{G'}, \mathcal{E}_\mu) = \ker(res^q), \tag{7.2}$$

which follows from considering the long exact sequence in sheaf-cohomology given by the pair $(\overline{S}_{G'}, \partial(\overline{S}_{G'}))$. We obtain

Lemma 7.3. *The $G'(\mathbb{A}_f)$-modules $H^q(S_{G'}, \mathcal{E}_\mu)$ and $H^q_!(S_{G'}, \mathcal{E}_\mu)$ are defined over $\mathbb{Q}(\mu)$.*

Proof. Both assertions follow from the fact that sheaf-cohomology can be computed using Betti-cohomology. To be more precise, we recall from Lem. 7.1 that E_μ is defined over its rationality field $\mathbb{Q}(\mu)$. Let $E_{\mu,\mathbb{Q}(\mu)}$ be a $\mathbb{Q}(\mu)$-structure on E_μ and let $H^q_B(S_{G'}, E_{\mu,\mathbb{Q}(\mu)})$ denote the Betti-cohomology of $S_{G'}$ with coefficients in $E_{\mu,\mathbb{Q}(\mu)}$. This is a $G'(\mathbb{A}_f)$-stable $\mathbb{Q}(\mu)$-vector space and defines a $\mathbb{Q}(\mu)$-structure:

$$H^q(S_{G'}, \mathcal{E}_\mu) \cong H^q_B(S_{G'}, E_{\mu,\mathbb{Q}(\mu)}) \otimes_{\mathbb{Q}(\mu)} \mathbb{C}.$$

Recalling (7.2), this also shows the analogous assertion for $H^q_!(S_{G'}, \mathcal{E}_\mu)$. □

Compare the previous lemma also with Clozel [8], p.122-123.

As another ingredient observe that for all $\sigma \in \mathrm{Aut}(\mathbb{C})$, there are natural σ-linear, $G'(\mathbb{A}_f)$-equivariant isomorphisms

$$\sigma^* : H^q(S_{G'}, \mathcal{E}_\mu) \to H^q(S_{G'}, {}^{\sigma}\mathcal{E}_\mu) \tag{7.4}$$

and

$$\sigma^*_! : H^q_!(S_{G'}, \mathcal{E}_\mu) \to H^q_!(S_{G'}, {}^{\sigma}\mathcal{E}_\mu). \tag{7.5}$$

7.3. **Combining automorphic forms and geometry.** Let E_μ be a highest weight representation of G'_∞ and let $\mathcal{J} = \mathcal{J}_{E_\mu}$ be the ideal of the center of the universal enveloping algebra of $\mathfrak{g}'_\mathbb{C}$, which annihilates the dual representation $E_\mu^\vee$ of E_μ. This ideal is of finite codimension and hence the subspace $\mathcal{A}_\mathcal{J}(G')$ of automorphic forms on $\mathbb{R}_+G'(F)\backslash G'(\mathbb{A})$, which are annihilated by some power of $\mathcal{J}$ is defined, cf. Section 3. Suppose furthermore that $E_\mu|_{\mathbb{R}_+} = 1$ in this subsection. Then the link between sheaf cohomology and automorphic forms is provided by

Theorem 7.6 (Franke [11], Thm. 18)**.** *There is an isomorphism of $G'(\mathbb{A}_f)$-modules*

$$H^q(S_{G'}, \mathcal{E}_\mu) \cong H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathcal{A}_\mathcal{J}(G') \otimes E_\mu).$$

Abbreviating

$$H^q_{\{P'\}}(G', E_\mu) := H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathcal{A}_{\mathcal{J},\{P'\}}(G') \otimes E_\mu)$$

and

$$H^q_{\{P'\},\varphi_{P'}}(G', E_\mu) := H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathcal{A}_{\mathcal{J},\{P'\},\varphi_{P'}}(G') \otimes E_\mu),$$

we obtain

Corollary 7.7. *The sheaf cohomology of $S_{G'}$ inherits from* (3.1) *and* (3.2) *a decomposition as $G'(\mathbb{A}_f)$-module:*

$$H^q(S_{G'}, \mathcal{E}_\mu) \cong \bigoplus_{\{P'\}} H^q_{\{P'\}}(G', E_\mu) \tag{7.8}$$

$$\cong \bigoplus_{\{P'\}} \bigoplus_{\varphi_{P'}} H^q_{\{P'\},\varphi_{P'}}(G', E_\mu). \tag{7.9}$$

7.4. Recall from Section 3 that the space $\mathcal{A}_{cusp,\mathcal{J}}(G')$, i.e., the subspace of all cuspidal automorphic forms in $\mathcal{A}_\mathcal{J}(G')$, can be identified with the summand $\mathcal{A}_{\mathcal{J},\{G'\}}(G')$ in (3.1). Hence, we call the summand in (7.8) index by $\{G'\}$ the space of *cuspidal cohomology* and denote it

$$\begin{aligned} H^q_{cusp}(G', E_\mu) &:= H^q_{\{G'\}}(G', E_\mu) \\ &= H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathcal{A}_{cusp,\mathcal{J}}(G') \otimes E_\mu). \end{aligned}$$

Now, let $\mathcal{A}_{dis,\mathcal{J}}(G')$ be the subspace of all square-integrable automorphic forms in $\mathcal{A}_\mathcal{J}(G')$. The quotient space $\mathbb{R}_+G'(F)\backslash G'(\mathbb{A})$ has finite volume hence $\mathcal{A}_{dis,\mathcal{J}}(G')$ is the space of all K'°_∞-finite, smooth functions $f \in L^2_{dis}(\mathbb{R}_+G'(F)\backslash G'(\mathbb{A}))$, which are annihilated by some power of $\mathcal{J}$. We will denote its cohomology by

$$H^q_{dis}(G', E_\mu) := H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathcal{A}_{dis,\mathcal{J}}(G') \otimes E_\mu).$$

Observe that by (7.8), there is a natural inclusion of $G'(\mathbb{A}_f)$-modules

$$H^q_{cusp}(G', E_\mu) \hookrightarrow H^q_{dis}(G', E_\mu).$$

Let us now refine this picture even more. If we fix a smooth character

$$\omega : \mathbb{R}_+Z'(F)\backslash Z'(\mathbb{A}) \to \mathbb{C}^*,$$

then the subspace of functions in $\mathcal{A}_{dis,\mathcal{J}}(G')$ (resp., $\mathcal{A}_{cusp,\mathcal{J}}(G')$), which have this given central character ω, decomposes as a direct Hilbert sum, the sum ranging over all (equivalence classes) of square-integrable (resp., cuspidal), irreducible automorphic representations with central character ω, which are annihilated by some power of $\mathcal{J}$:

$$\mathcal{A}_{dis,\mathcal{J}}(G', \omega) \cong \widehat{\bigoplus}_{\Pi' \text{ square-int.}} \Pi', \tag{7.10}$$

resp.,

$$\mathcal{A}_{cusp,\mathcal{J}}(G', \omega) \cong \widehat{\bigoplus}_{\Pi' \text{ cuspidal.}} \Pi'. \tag{7.11}$$

Once more, these spaces define in a natural way $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-modules and if E_μ is a highest weight representation of G'_∞, we define

$$H^q_{dis,\omega}(G', E_\mu) := H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathcal{A}_{dis,\mathcal{J}}(G',\omega) \otimes E_\mu) \tag{7.12}$$

$$\cong \bigoplus_{\Pi' \text{ square-int.}} H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \Pi'_\infty \otimes E_\mu) \otimes \Pi'_f \tag{7.13}$$

and

$$H^q_{cusp,\omega}(G', E_\mu) := H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathcal{A}_{cusp,\mathcal{J}}(G',\omega) \otimes E_\mu) \tag{7.14}$$

$$\cong \bigoplus_{\Pi' \text{ cuspidal}} H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \Pi'_\infty \otimes E_\mu) \otimes \Pi'_f. \tag{7.15}$$

Both spaces are $G'(\mathbb{A}_f)$-modules and their decompositions are inherited by (7.10), resp., (7.11).

By Thm. 7.6 it is justified to talk about the image of all cohomology spaces constructed in Section 7.3 and 7.4 in $H^q(S_{G'}, \mathcal{E}_\mu)$. Let us denote these various images by overlining "H". In particular, this applies to $H^q_{cusp}(G', E_\mu)$ and $H^q_{dis}(G', E_\mu)$ (resp., $H^q_{cusp,\omega}(G', E_\mu)$ and $H^q_{dis,\omega}(G', E_\mu)$), so, e.g., $\mathrm{Im} H^q_{cusp}(G', E_\mu) =: \bar{H}^q_{cusp}(G', E_\mu)$. Then Cor. 7.7 implies the following proposition, which describes the interplay between interior cohomology and the various cohomology spaces constructed in this section.

Proposition 7.16. *There is the following commutative diagram of $G'(\mathbb{A}_f)$-modules.*

$$\begin{tikzcd}
H^q_{cusp}(G', E_\mu) \arrow[r, "\cong"] & \bar{H}^q_{cusp}(G', E_\mu) \arrow[r, hook] & H^q_!(S_{G'}, \mathcal{E}_\mu) \arrow[r, hook] & \bar{H}^q_{dis}(G', E_\mu) \\
H^q_{cusp,\omega}(G', E_\mu) \arrow[r, "\cong"] & \bar{H}^q_{cusp,\omega}(G', E_\mu) \arrow[u, hook] \arrow[rr, hook] & & \bar{H}^q_{dis,\omega}(G', E_\mu) \arrow[u, hook]
\end{tikzcd}$$

7.5. **Three results in the split case.** Having made some preparatory work in Sections 7.1–7.4, let us now recall one of the main results in Section 3 of Clozel [8]:

Theorem 7.17 ([8], Thm. 3.13)**.** *Let $\Pi = \Pi_\infty \otimes \Pi_f \in \mathscr{D}(G)$ be regular algebraic and cuspidal. Then for all $\sigma \in \mathrm{Aut}(\mathbb{C})$, ${}^\sigma\Pi_f$ is the finite part of a regular algebraic, cuspidal representation $\Xi \in \mathscr{D}(G)$ such that ${}^\sigma\Pi_\infty \cong \Xi_\infty$.*

We may reformulate Thm. 7.17 within the setup developed in Sections 7.2-7.4 applied to the special case $D = F$, i.e., $G' = G$. As Π is regular algebraic and cuspidal, Π_∞ is cohomological with respect to some highest weight representation E_μ of G_∞, cf. Thm. 6.3. Therefore, by (7.15), Π_f appears as an irreducible subspace of $H^q_{cusp,\omega}(G, E_\mu)$ for some degree q and ω being the central character of Π (where we can assume without loss of generality that $\omega|_{\mathbb{R}_+} = 1$). Hence, by Prop. 7.16, Π_f is even an irreducible submodule of $H^q_{cusp}(G, E_\mu)$. In view of the decomposition of the cohomology of S_G, cf. Cor. 7.7, one may hence reformulate Thm. 7.17 as follows:

For any highest weight representation E_μ and any automorphism $\sigma \in \mathrm{Aut}(\mathbb{C})$, the summand $H^q_{\{G\}}(G, E_\mu)$ of $H^q(S_G, \mathcal{E}_\mu)$, being cuspidal cohomology, is mapped by σ^* (cf. 7.4) isomorphically onto the summand $H^q_{\{G\}}(G, {}^\sigma E_\mu)$ of $H^q(S_G, {}^\sigma\mathcal{E}_\mu)$

It is now clear why the following theorem of Franke is a generalization of Thm. 7.17:

Theorem 7.18 ([11], Thm. 20)**.** *Let E_μ be a highest weight representation of G_∞ and $\sigma \in \mathrm{Aut}(\mathbb{C})$ an automorphism. For each associate class of parabolic F-subgroups $\{P\}$ the summand $H^q_{\{P\}}(G, E_\mu)$ of $H^q(S_G, \mathcal{E}_\mu)$, is mapped by σ^* isomorphically onto the corresponding summand $H^q_{\{P\}}(G, {}^\sigma E_\mu)$ of $H^q(S_G, {}^\sigma\mathcal{E}_\mu)$. In other words, σ^* respects the decomposition of sheaf cohomology along associate classes of parabolic F-subgroups.*

This latter theorem was refined further by Franke–Schwermer in [12]:

Theorem 7.19 ([12], Thm. 4.3). *Let E_μ be a highest weight representation of G_∞ and $\sigma \in \mathrm{Aut}(\mathbb{C})$ an automorphism. For each associate class of parabolic F-subgroups $\{P\}$, and each associate class of cuspidal automorphic representations φ_P, the summand $H^q_{\{P\},\varphi_P}(G, E_\mu)$ of $H^q(S_G, \mathcal{E}_\mu)$, is mapped by σ^* isomorphically onto the summand $H^q_{\{P\},{}^\sigma\varphi_P}(G, {}^\sigma E_\mu)$ of $H^q(S_G, {}^\sigma\mathcal{E}_\mu)$ for a unique associate class ${}^\sigma\varphi_P$.*

We devote the next two subsections to showing certain generalizations of the above theorems to G'.

7.6. **A generalization of Theorems 7.17, 7.18 and 7.19.**

Proposition 7.20. *Let $\Pi' \in \mathscr{D}(G')$ be cuspidal and cohomological with respect to a highest weight representation E_μ. Then for all $\sigma \in \mathrm{Aut}(\mathbb{C})$, ${}^\sigma\Pi'_f$ is the finite part of a discrete series representation $\Xi' \in \mathscr{D}(G')$, which is cohomological with respect to ${}^\sigma E_\mu$. If E_μ is regular, then Ξ' is cuspidal.*

Proof. By assumption Π'_∞ is cohomological with respect to the highest weight representation E_μ. In particular, by [7], I, Thm. 5.3.(ii), the central character $\omega_{\Pi'_\infty}$ of Π'_∞ and the central character of $E^\vee_\mu$ coincide. Therefore, we may henceforth assume without loss of generality that the central characters of Π'_∞ and E_μ are both trivial on $\mathbb{R}_+$.

To proceed, let b be a degree in which the $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-cohomology of $\Pi'_\infty \otimes E_\mu$ does not vanish. Then $H^b(\mathfrak{g}'_\infty, K'^\circ_\infty, \Pi'_\infty \otimes E_\mu)$ is a non-trivial, but finite-dimensional $\mathbb{C}$-vector space on which $G'(\mathbb{A}_f)$ acts trivially. We may therefore embed

$$\Pi'_f \hookrightarrow H^b(\mathfrak{g}'_\infty, K'^\circ_\infty, \Pi'_\infty \otimes E_\mu) \otimes \Pi'_f.$$

According to (7.15), this implies that Π'_f appears as an irreducible $G'(\mathbb{A}_f)$-submodule in $H^b_{cusp,\omega}(G', E_\mu)$, with ω the central character of Π'. So, by Prop. 7.16, we may view Π'_f as an irreducible $G'(\mathbb{A}_f)$-submodule of $H^b_!(S_{G'}, \mathcal{E}_\mu)$. Recall from (7.5) that for each $\sigma \in \mathrm{Aut}(\mathbb{C})$ there is a σ-linear, $G'(\mathbb{A}_f)$-equivariant isomorphism

$$\sigma^*_! : H^b_!(S_{G'}, \mathcal{E}_\mu) \longrightarrow H^b_!(S_{G'}, {}^\sigma\mathcal{E}_\mu).$$

Hence, ${}^\sigma\Pi'_f$ appears as an irreducible submodule of $H^b_!(S_{G'}, {}^\sigma\mathcal{E}_\mu)$ and using Prop. 7.16 again, even as an irreducible submodule of $\bar{H}^b_{dis}(G', {}^\sigma E_\mu)$. By (7.10), there must hence be a discrete series representation $\Xi' \in \mathscr{D}(G')$, such that Ξ'_∞ is cohomological with respect to ${}^\sigma E_\mu$ and $\Xi'_f \cong {}^\sigma\Pi'_f$. This proves the first claim.

For the second claim, assume E_μ to be regular. Then ${}^\sigma E_\mu$ is regular, too. It is well–known that under these conditions each local archimedean component of Ξ'_∞ must be essentially tempered. This follows from the Vogan–Zuckerman's condition [40] (5.1), p. 73, (which is (4.8) here) together with the last paragraph on p. 58, *ibidem*; see also Li–Schwermer [24], Prop. 4.2, or Franke [11], p. 258. Now, Wallach [42], Thm. 4.3, shows that Ξ' is cuspidal. ☐

Remark 7.21. As it is obvious from the proof of Prop. 7.20, Ξ' will also be cuspidal under the weaker condition that E_μ only admits essentially tempered irreducible admissible representations of G'_∞, which have non–vanishing $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-cohomology twisted by E_μ. For an example of a non–regular coefficient module E_μ, which has this property, see Grobner [14], Prop. 3.5.

Theorem 7.22. *Let E_μ be a regular highest weight representation of G'_∞ and let $\sigma \in \mathrm{Aut}(\mathbb{C})$. For each associate class of parabolic F-subgroups $\{P'\}$, and each associate class of cuspidal automorphic representations $\varphi_{P'}$, the summand $H^q_{\{P'\},\varphi_{P'}}(G', E_\mu)$ of $H^q(S_{G'}, \mathcal{E}_\mu)$, is mapped by σ^* isomorphically onto the summand $H^q_{\{P'\},{}^\sigma\varphi_{P'}}(G', {}^\sigma E_\mu)$ of $H^q(S_{G'}, {}^\sigma\mathcal{E}_\mu)$ for a unique associate class ${}^\sigma\varphi_{P'}$:*

$$H^q_{\{P'\},\varphi_{P'}}(G', E_\mu) \xrightarrow[\sigma*]{\cong} H^q_{\{P'\},{}^\sigma\varphi_{P'}}(G', {}^\sigma E_\mu).$$

Assuming $H^q_{\{P'\},\varphi_{P'}}(G', E_\mu) \neq 0$ and letting $\{P'\}$ be represented by the parabolic F-subgroup P' with Levi factor L' and $\varphi_{P'}$ be represented by the cuspidal automorphic representation Π' of $L'(\mathbb{A})$, then $\Pi' \otimes \rho_{P'}$ is cohomological. The σ-twist of its finite part $\Pi'_f \otimes \rho_{P'_f}$ is the finite part of a representation Ξ' of $L'(\mathbb{A})$, which is indeed cuspidal and the associate class ${}^\sigma\varphi_{P'}$ is uniquely determined by being represented by the representation $\Xi' \otimes \rho^{-1}_{P'}$.

Proof. Recall that $G' = GL'_m$ for some $m \geq 1$. Let $\{P'\}$ be the associate class of the parabolic F-subgroup P' with Levi factor $L' = \prod_{i=1}^{r} GL'_{m_i}$, $\sum_{i=1}^{r} m_i = m$ and $\varphi_{P'}$ be represented by the cuspidal automorphic representation Π' of $L'(\mathbb{A})$. Assume that $H^q_{\{P'\},\varphi_{P'}}(G', E_\mu) \neq 0$.

If $r = 1$, i.e., $P' = G'$, each associate class $\varphi_{G'}$ of cuspidal automorphic representations is a singleton consisting of an equivalence class of a cuspidal automorphic representation Π' of $G'(\mathbb{A})$. Hence, if $H^q_{\{G'\},\varphi_{G'}}(G', E_\mu) \neq 0$, then all its irreducible $G'(\mathbb{A}_f)$-subquotients are in fact irreducible subrepresentations isomorphic to the finite part of the cohomological, cuspidal automorphic representation Π', cf. Section 7.4. So σ^* maps such a $G'(\mathbb{A}_f)$-subquotient onto the $G'(\mathbb{A}_f)$-representation ${}^\sigma\Pi'_f$. According to Prop. 7.20, ${}^\sigma\Pi'_f$ is the finite part of a discrete series representation Ξ' which has non-zero cohomology with respect to ${}^\sigma E_\mu$. As E_μ is regular, Ξ' is cuspidal by Prop. 7.20 and the assertion of the theorem follows in this case.

Now, let $r \neq 1$, i.e., $P' \neq G'$. By the very construction of the space $\mathcal{A}_{\mathcal{J},\{P'\},\varphi_{P'}}(G')$, cf. Section 3, and the regularity of E_μ, there is an isomorphism of $G'(\mathbb{A}_f)$-modules

$$H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathrm{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\tilde{\Pi}' \otimes S(\mathfrak{a}^{G'}_{P',\mathbb{C}})] \otimes E_\mu) \xrightarrow{\sim} H^q_{\{P'\},\varphi_{P'}}(G', E_\mu),$$

where $\tilde{\Pi}'$ is as in Section 3 and $S(\mathfrak{a}^{G'}_{P',\mathbb{C}})$ is the algebra of differential operators on the finite-dimensional complex space $\mathfrak{a}^{G'}_{P',\mathbb{C}}$ of variables of the Eisenstein series attached to $\tilde{\Pi}'$. All relevant details concerning this construction are contained in Franke–Schwermer [12] 1.2-1.4, 3.3 to which we refer. The existence of the above isomorphism is proved in Franke [11] Thm. 19 II. Let $\mathfrak{m}'_\infty$ be the reductive Lie algebra obtained from $\mathfrak{l}'_\infty$ by quotienting out the diagonally embedded commutative Lie algebra $\mathfrak{a}^{G'}_{P'} \hookrightarrow \mathfrak{l}'_\infty$ and $K'_{L',\infty}$ be the intersection $K'^\circ_\infty \cap L'_\infty$. Then, in [11], pp. 256-257, it is shown that as a $G'(\mathbb{A}_f)$-module,

$$H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathrm{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\tilde{\Pi}' \otimes S(\mathfrak{a}^{G'}_{P',\mathbb{C}})] \otimes E_\mu) \cong \tag{7.23}$$

$$\mathrm{Ind}_{P'(\mathbb{A}_f)}^{G'(\mathbb{A}_f)}\left[H^{q-l}(\mathfrak{m}'_\infty, K'_{L',\infty}, \tilde{\Pi}'_\infty \otimes E_{\mu_w,\mathfrak{m}'_\infty}) \otimes \Pi'_f\right],$$

where $E_{\mu_w,\mathfrak{m}'_\infty}$ is a uniquely determined highest weight representation of $\mathfrak{m}'_\infty$ and l is a certain shift in degrees. In fact, $E_{\mu_w,\mathfrak{m}'_\infty}$ is the restriction of the irreducible L'_∞-representation E_{μ_w} of highest weight $\mu_w = w(\mu + \rho) - \rho$, w being a uniquely determined Kostant representative in $W^{P'}$, cf. Borel–Wallach [7] III 1.4 and III 3.3 for a definition of $W^{P'}$ and the uniqueness of w. Then, the shift in degrees l equals the length $l(w)$ of w, see again [7] III 3.3. On the other hand, directly by [7], III Thm. 3.3, one obtains that

$$H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathrm{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\Pi'] \otimes E_\mu) \cong \tag{7.24}$$

$$\bigoplus_{a+b=q-l} \mathrm{Ind}_{P'(\mathbb{A}_f)}^{G'(\mathbb{A}_f)}\left[H^a(\mathfrak{m}'_\infty, K'_{L',\infty}, \tilde{\Pi}'_\infty \otimes E_{\mu_w,\mathfrak{m}'_\infty}) \otimes \bigwedge^b \check{a}^{G'}_{P',\mathbb{C}} \otimes \Pi'_f\right],$$

revealing (7.23) as a direct summand in (7.24) attached to $b = 0$. As a consequence, there is also a surjective $G'(\mathbb{A}_f)$-module homomorphism

$$H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathrm{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\Pi'] \otimes E_\mu) \twoheadrightarrow H^q_{\{P'\},\varphi_{P'}}(G', E_\mu). \tag{7.25}$$

Now, invoking [7] III 3.4.(5) and III 3.4.(14), we see that $H^{q-l}(\mathfrak{l}'_\infty, K'_{L',\infty}, \Pi'_\infty \otimes \rho_{P'_\infty} \otimes E_{\mu_w}) \neq 0$, where $\rho_{P'_\infty}$ is the archimedean component of the character $\rho_{P'} = \rho_{P'_\infty} \otimes \rho_{P'_f}$, which is defined as usual via the Harish-Chandra homomorphism. Observe that the group $K'_{L',\infty}$ is in general neither connected nor contains the full connected component $Z'^\circ_{L',\infty}$ of the center of L'_∞. However, a short moment of thought, using [7] I 5.1 and Thm. I 5.3, shows that $\Pi'_\infty \otimes \rho_{P'_\infty}$ is really cohomological in our sense, made precise in Sect. 2.4, with respect to the irreducible, algebraic representation E_{μ_w} of L'_∞. One may check that E_μ being regular implies that also E_{μ_w} is regular. Therefore, we may apply Prop. 7.20 to the cuspidal automorphic representation $\Pi' \otimes \rho_{P'}$ and obtain a cuspidal automorphic representation Ξ' of $L'(\mathbb{A})$ with the property that

$\Xi'_f \cong {}^\sigma\Pi'_f \otimes {}^\sigma\rho_{P'_f}$. By Strong Multiplicity One and Multiplicity One for discrete series representations of $L'(\mathbb{A})$, we obtain furthermore that

$$H^b(\mathfrak{l}'_\infty, (K'_{L',\infty}Z'_{L',\infty})^\circ, \Pi'_\infty \otimes \rho_{P'_\infty} \otimes E_{\mu_w}) \cong H^b(\mathfrak{l}'_\infty, (K'_{L',\infty}Z'_{L',\infty})^\circ, \Xi'_\infty \otimes {}^\sigma E_{\mu_w}),$$

for all degrees b. From this one may also derive that

$$H^b(\mathfrak{l}'_\infty, (K'_{L',\infty})^\circ, \Pi'_\infty \otimes \rho_{P'_\infty} \otimes E_{\mu_w}) \cong H^b(\mathfrak{l}'_\infty, (K'_{L',\infty})^\circ, \Xi'_\infty \otimes {}^\sigma E_{\mu_w}) \tag{7.26}$$

holds for all degrees b. Keeping this in mind, we let $\partial_{P'}S_{G'} := P'(F)\backslash G'(\mathbb{A})/K'^\circ_\infty$. It is known that

$$H^q(\partial_{P'}S_{G'}, \mathcal{E}_\mu) \cong {}^a\mathrm{Ind}_{\pi_0(P'_\infty)\times P'(\mathbb{A}_f)}^{\pi_0(G'_\infty)\times G'(\mathbb{A}_f)}\left[\bigoplus_{w\in W^{P'}} H^{q-l(w)}(L'(F)\backslash L'(\mathbb{A})/K'_{L',\infty}, \mathcal{E}_{\mu_w})\right],$$

"^{a}Ind" denoting un-normalized or algebraic induction, cf. [33], 7.1–7.2. This isomorphism allows one to define the $G'(\mathbb{A}_f)$- submodule

$$H^q_{cusp}(\partial_{P'}S_{G'}, \mathcal{E}_\mu) := {}^a\mathrm{Ind}_{\pi_0(P'_\infty)\times P'(\mathbb{A}_f)}^{\pi_0(G'_\infty)\times G'(\mathbb{A}_f)}\left[\bigoplus_{w\in W^{P'}} \bigoplus_{\Phi'} H^{q-l(w)}(\mathfrak{l}'_\infty, K'_{L',\infty}, \Phi'_\infty \otimes E_{\mu_w}) \otimes \Phi'_f\right],$$

the second sum ranging over all equivalence classes of cuspidal automorphic representations Φ' of $L'(\mathbb{A})$ which satisfy $\omega_{\Phi'_\infty}|_{Z'^\circ_{L',\infty}} = \omega^{-1}_{E_{\mu_w}}|_{Z'^\circ_{L',\infty}}$. It is an easy observation that the kernel of the surjective map $\pi_0(P'_\infty) \to \pi_0(G'_\infty)$ is equal to the image of $K'_{L',\infty}/(K'_{L',\infty})^\circ$ in $\pi_0(P'_\infty)$. Hence, one may rewrite the above by

$$\begin{aligned}\mathrm{Ind}_{P'(\mathbb{A}_f)}^{G'(\mathbb{A}_f)}&\left[\bigoplus_{w\in W^{P'}} \bigoplus_{\Phi'} H^{q-l(w)}(\mathfrak{l}'_\infty, K'_{L',\infty}, \Phi'_\infty \otimes \rho_{P'_\infty} \otimes E_{\mu_w}) \otimes \Phi'_f\right]\\ &\cong \bigoplus_{\Phi'} H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathrm{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\Phi'] \otimes E_\mu),\end{aligned}$$

where the Φ' now range over all equivalence classes of cuspidal automorphic representations of $L'(\mathbb{A})$ which satisfy the modified condition $\omega_{\Phi'_\infty}|_{Z'^\circ_{L',\infty}} = \left(\omega_{E_{\mu_w}} \cdot \rho_{P'_\infty}\right)^{-1}|_{Z'^\circ_{L',\infty}}$ for the corresponding, uniquely determined Kostant representative $w \in W^{P'}$. In particular, by what we said above, $H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathrm{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\Pi'] \otimes E_\mu)$ is a $G'(\mathbb{A}_f)$-submodule of $H^q_{cusp}(\partial_{P'}S_{G'}, \mathcal{E}_\mu)$ and hence of the sheaf cohomology $H^q(\partial_{P'}S_{G'}, \mathcal{E}_\mu)$. Whence, it makes sense to apply σ^* to the latter $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-cohomology space. Choosing ${}^\sigma\varphi_{P'}$ to be the associate class which is represented by $\Xi' \otimes \rho^{-1}_{P'}$, we obtain the following diagram

$$\begin{tikzcd}
H^q(S_{G'}, \mathcal{E}_\mu) \arrow[d, "\sigma^*"] & H^q_{\{P'\},\varphi_{P'}}(G', E_\mu) \arrow[l, hook] & H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathrm{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\Pi'] \otimes E_\mu) \arrow[l, two heads] \arrow[r, hook] & H^q(\partial_{P'}S_{G'}, \mathcal{E}_\mu) \arrow[d, "\sigma^*"] \\
H^q(S_{G'}, {}^\sigma\mathcal{E}_\mu) & H^q_{\{P'\},{}^\sigma\varphi_{P'}}(G', {}^\sigma E_\mu) \arrow[l, hook] & H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathrm{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\Xi' \otimes \rho^{-1}_{P'}] \otimes {}^\sigma E_\mu) \arrow[l, two heads] \arrow[r, hook] & H^q(\partial_{P'}S_{G'}, {}^\sigma\mathcal{E}_\mu)
\end{tikzcd}$$

Equation (7.26) together with Strong Multiplicity One and Multiplicity One for discrete series representations of $L'(\mathbb{A})$ finally show that the restriction of $\sigma^* : H^q(\partial_{P'}S_{G'}, \mathcal{E}_\mu) \to H^q(\partial_{P'}S_{G'}, {}^\sigma\mathcal{E}_\mu)$ to the space $H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathrm{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\Pi'] \otimes E_\mu)$ provides us with a commutative diagram

$$\begin{tikzcd}
H^q(S_{G'}, \mathcal{E}_\mu) \arrow[d, "\sigma^*"] & H^q_{\{P'\},\varphi_{P'}}(G', E_\mu) \arrow[l, hook] & H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathrm{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\Pi'] \otimes E_\mu) \arrow[l, two heads] \arrow[d, "\sigma^*"] \\
H^q(S_{G'}, {}^\sigma\mathcal{E}_\mu) & H^q_{\{P'\},{}^\sigma\varphi_{P'}}(G', {}^\sigma E_\mu) \arrow[l, hook] & H^q(\mathfrak{g}'_\infty, K'^\circ_\infty, \mathrm{Ind}_{P'(\mathbb{A})}^{G'(\mathbb{A})}[\Xi' \otimes \rho^{-1}_{P'}] \otimes {}^\sigma E_\mu) \arrow[l, two heads]
\end{tikzcd}$$

This settles the case $r \neq 1$ and hence the theorem. □

7.7. **A complementary view on Thm. 7.17.** Let now $\Pi' \in \mathscr{D}(G')$ be regular algebraic and assume that its global Jacquet-Langlands transfer $JL(\Pi')$ is cuspidal - the standard assumption we had to make in Section 6, in order to obtain proper generalizations of the various theorems on regular algebraic representations in the split case. The next theorem is complementary to Thm. 7.22 and shows that for any automorphism $\sigma \in \mathrm{Aut}(\mathbb{C})$, the $G'(\mathbb{A}_f)$-equivariant isomorphism σ^* respects these properties. Hence, it may also be seen as another generalization of Thm. 7.17.

Theorem 7.27. *Let $\Pi' \in \mathscr{D}(G')$ be regular algebraic and assume that $JL(\Pi')$ is cuspidal. Then for all $\sigma \in \mathrm{Aut}(\mathbb{C})$, ${}^\sigma\Pi'_f$ is the finite part of a cuspidal representation $\Xi' \in \mathscr{D}(G')$ which is regular algebraic and $JL(\Xi')$ is cuspidal.*

Proof. Let $\Pi' = \Pi'_\infty \otimes \Pi'_f \in \mathscr{D}(G')$ be as in the statement of the theorem. By Thm. 6.9 we know that Π'_∞ is cohomological and by Cor. 5.5 that Π' is cuspidal. Hence, by Prop. 7.20, there is a discrete series representation Ξ' of $G'(\mathbb{A})$, which satisfies $\Xi'_f \cong {}^\sigma\Pi'_f$ and has non–vanishing $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-cohomology with respect to ${}^\sigma E_\mu$. So we can finish the proof by showing that $JL(\Xi')$ is regular algebraic and cuspidal.

In order to do so, recall from Thm. 7.17 that ${}^\sigma JL(\Pi')_f$ is the finite part of a regular algebraic, cuspidal representation Ξ of $G(\mathbb{A})$. Now we observe that at all split places $v \in V_f$

$$JL(\Xi')_v \cong \Xi'_v \cong ({}^\sigma\Pi')_v \cong {}^\sigma(\Pi'_v) \cong {}^\sigma JL(\Pi')_v \cong \Xi_v.$$

Therefore, the discrete series representation $JL(\Xi')$ and the cuspidal representation Ξ are isomorphic almost everywhere, and hence by the Strong Multiplicity One Theorem and the Multiplicity One Theorem for discrete series representations of $G(\mathbb{A})$,

$$JL(\Xi') = \Xi.$$

In particular, $JL(\Xi')$ is cuspidal and regular algebraic, whence so is Ξ'. □

Remark 7.28. Our Thm. 7.27 is also a proper generalization of Waldspurger's Thm. II.3.2 in [41].

8. Results on rationality fields

8.1. The next theorem generalizes a result which is well–known in the split case (cf., e.g., Shimura [35], Harder [16] p.80, Waldspurger [41], Cor. I.8.3 and first line of p. 153, and Clozel [8]).

Theorem 8.1. *Let $\Pi' \in \mathscr{D}(G')$ be regular algebraic and assume that $JL(\Pi')$ is cuspidal. Then $\mathbb{Q}(\Pi'_f)$ is a number field.*

Proof. Let Π' be regular algebraic and assume that $JL(\Pi')$ is cuspidal. By Thm. 7.27, for any automorphism $\sigma \in \mathrm{Aut}(\mathbb{C})$, ${}^\sigma\Pi'_f$ is the finite part of a cuspidal automorphic representation Ξ' which has non–vanishing cohomology with respect to ${}^\sigma E_\mu$. Using Prop. 7.16 we hence obtain inclusions

$${}^\sigma\Pi'_f \hookrightarrow H^q_{cusp}(G', {}^\sigma E_\mu),$$

for all $\sigma \in \mathrm{Aut}(\mathbb{C})$. Take an open, compact subgroup K'_f of $G'(\mathbb{A}_f)$ which is small enough such that ${\Pi'_f}^{K'_f} \neq 0$. Then for each σ, $H^q_{cusp}(G', {}^\sigma E_\mu)^{K'_f}$ is finite-dimensional. Therefore, $|\mathrm{Aut}(\mathbb{C})/\mathfrak{S}(\Pi'_f)|$ is finite. By [19], V Lem. 2.9, $|\mathrm{Aut}(\mathbb{C})/\mathfrak{S}(\Pi'_f)|$ is an upper bound for $\dim_{\mathbb{Q}}(\mathbb{Q}(\Pi'_f)) = [\mathbb{Q}(\Pi'_f) : \mathbb{Q}]$. So, $\mathbb{Q}(\Pi'_f)$ is a number field. □

8.2. Let us also discuss how the local rationality fields $\mathbb{Q}(\Pi'_v)$, $v \in V_f$, and $\mathbb{Q}(\Pi'_f)$ are connected to each other.

Proposition 8.2. *Let $\Pi' \in \mathscr{D}(G')$ be regular algebraic and assume that $JL(\Pi')$ is cuspidal. Then $\mathbb{Q}(\Pi'_f)$ is the compositum of the fields $\mathbb{Q}(\Pi'_v)$, $v \in V_f - S$, $S \subset V_f$ an arbitrary finite subset containing all non-archimedean places where Π'_v ramifies.*

The proof of Prop. 8.2 requires a few preparatory lemmas. Let $v \in V_f$ be a split, non-archimedean place. So $G'_v = G_v = GL_n(F_v)$ and let ϖ_v be an uniformizer for the maximal ideal of the ring of integers $\mathcal{O}_v$ of F_v. If Π_v is an unramified, irreducible admissible representation of G'_v, then it is well–known that

$$\Pi_v \cong \mathrm{Ind}_{B_v}^{G_v}[\chi_1 \otimes ... \otimes \chi_n],$$

for some unramified characters $\chi_j : F_v^* \to \mathbb{C}^*$ and B_v being the Borel subgroup of G_v. We put

$$\alpha_j^{\Pi'_v} := |\varpi_v|_v^{\frac{n-1}{2}} \chi_j(\varpi_v),$$

with $|\cdot|_v$ being the normalized absolute value on F_v. Furthermore, for any automorphism $\sigma \in \mathrm{Aut}(\mathbb{C})$ denote by

$$\epsilon_{\sigma,v} = \frac{|\det'|_v^{1/2}}{\sigma(|\det'|_v^{1/2})}.$$

Then $\epsilon_{\sigma,v}$ is a quadratic character, because by the very definition $|\cdot|_v$ takes values in the rational numbers, whence $|\det'|_v = \sigma(|\det'|_v)$ for any automorphism $\sigma \in \mathrm{Aut}(\mathbb{C})$. As a last ingredient, recall the *elementary symmetric polynomials* $f_1, ..., f_n$, cf., e.g., [19] Appendix to Chp. V:

$$f_j(x_1, ..., x_n) = \sum_{1 \leq i_1 < ... < i_j \leq n} x_{i_1} x_{i_2} ... x_{i_j}.$$

The next lemma is a generalization of Waldspurger [41], Lem. I.2.3.

Lemma 8.3. *Let $v \in V_f$ be a split, non-archimedean place and Π_v an unramified, irreducible admissible representation of $G'_v = G_v = GL_n(F_v)$. Then*

$$\mathbb{Q}(\Pi_v) = \mathbb{Q}\left(f_1(\alpha_1^{\Pi'_v}, ..., \alpha_n^{\Pi'_v}), ..., f_n(\alpha_1^{\Pi'_v}, ..., \alpha_n^{\Pi'_v})\right).$$

Proof.

$$\begin{array}{lcl}
\sigma \in \mathfrak{S}(\Pi_v) & \Leftrightarrow & {}^\sigma\Pi_v \cong \Pi_v \\
 & \Leftrightarrow & {}^\sigma\mathrm{Ind}_{B_v}^{G_v}[\chi_1 \otimes ... \otimes \chi_n] \cong \mathrm{Ind}_{B_v}^{G_v}[\chi_1 \otimes ... \otimes \chi_n] \\
 & \underset{\text{[8], Lem. 3.5.(ii)}}{\Leftrightarrow} & \mathrm{Ind}_{B_v}^{G_v}[{}^\sigma\chi_1 \epsilon_{\sigma,v}^{n-1} \otimes ... \otimes {}^\sigma\chi_n \epsilon_{\sigma,v}^{n-1}] \cong \mathrm{Ind}_{B_v}^{G_v}[\chi_1 \otimes ... \otimes \chi_n] \\
 & \Leftrightarrow & \{{}^\sigma\chi_1 \epsilon_{\sigma,v}^{n-1}, ..., {}^\sigma\chi_n \epsilon_{\sigma,v}^{n-1}\} = \{\chi_1, ..., \chi_n\} \\
 & \Leftrightarrow & \sigma\left(f_j(\alpha_1^{\Pi'_v}, ..., \alpha_n^{\Pi'_v})\right) = f_j(\alpha_1^{\Pi'_v}, ..., \alpha_n^{\Pi'_v}) \quad \forall 1 \leq j \leq n \\
 & \Leftrightarrow & \sigma \in \mathrm{Aut}\left(\mathbb{C}\big/\mathbb{Q}\left(f_1(\alpha_1^{\Pi'_v}, ..., \alpha_n^{\Pi'_v}), ..., f_n(\alpha_1^{\Pi'_v}, ..., \alpha_n^{\Pi'_v})\right)\right).
\end{array}$$

Taking fixed fields proves the lemma. □

Lemma 8.4. *Let $v \in V_f$ be a split, non-archimedean place and Π_v an unramified, irreducible admissible representation of $G'_v = G_v = GL_n(F_v)$. Assume that $\mathbb{Q}(\Pi_v)$ is a number field. Then for any finite Galois extension $\mathbb{F}_v$ of $\mathbb{Q}$, containing $\mathbb{Q}(\Pi_v)$,*

$$\mathrm{Gal}(\mathbb{F}_v/\mathbb{Q}(\Pi_v)) = \{\sigma \in \mathrm{Gal}(\mathbb{F}_v/\mathbb{Q}) | {}^{\tilde\sigma}\Pi_v \cong \Pi_v \textit{ for all lifts } \tilde\sigma \in \mathrm{Aut}(\mathbb{C})\}.$$

Moreover, the restriction map $\mathrm{Aut}(\mathbb{C}) \to \mathrm{Gal}(\mathbb{F}_v/\mathbb{Q})$, sending τ to $\tau|_{\mathbb{F}_v}$, maps $\mathfrak{S}(\Pi_v)$ onto $\mathrm{Gal}(\mathbb{F}_v/\mathbb{Q}(\Pi_v))$.

Proof. By Lem. 8.3,

$${}^{\tilde\sigma}\Pi_v \cong \Pi_v \Leftrightarrow \tilde\sigma\left(f_j(\alpha_1^{\Pi'_v}, ..., \alpha_n^{\Pi'_v})\right) = f_j(\alpha_1^{\Pi'_v}, ..., \alpha_n^{\Pi'_v}) \quad \forall 1 \leq j \leq n.$$

But since $f_j(\alpha_1^{\Pi'_v}, ..., \alpha_n^{\Pi'_v}) \in \mathbb{Q}(\Pi_v) \subseteq \mathbb{F}_v$ for all $1 \leq j \leq n$, we obtain

$$\mathrm{Gal}(\mathbb{F}_v/\mathbb{Q}(\Pi_v)) = \{\sigma \in \mathrm{Gal}(\mathbb{F}_v/\mathbb{Q}) | {}^{\tilde\sigma}\Pi_v \cong \Pi_v \text{ for all lifts } \tilde\sigma \in \mathrm{Aut}(\mathbb{C})\}.$$

The last assertion is obvious. □

Lemma 8.5. *Let $\Pi' \in \mathscr{D}(G')$ be regular algebraic and assume that $JL(\Pi')$ is cuspidal and let $S \subset V_f$ be an arbitrary finite set containing all non-archimedean places where Π'_v ramifies. Denote by $\mathbb{E}$ the compositum of the fields $\mathbb{Q}(\Pi'_v)$, $v \in V_f - S$. Then for any finite Galois extension $\mathbb{F}$ of $\mathbb{Q}$, containing $\mathbb{Q}(\Pi'_f)$, the restriction map $r_{\mathbb{F}} : \mathrm{Aut}(\mathbb{C}) \to \mathrm{Gal}(\mathbb{F}/\mathbb{Q})$, sending τ to $\tau|_{\mathbb{F}}$, maps $\mathfrak{S}(\Pi'_f)$ onto $\mathrm{Gal}(\mathbb{F}/\mathbb{E})$.*

Proof. Recall from Thm. 8.1 that under the present assumptions $\mathbb{Q}(\Pi'_f)$ is a number field. So it makes sense to talk about a finite Galois extension $\mathbb{F}$ of $\mathbb{Q}$, containing $\mathbb{Q}(\Pi'_f)$. We get

$$\begin{aligned}
\mathrm{Gal}(\mathbb{F}/\mathbb{E}) &= \bigcap_{v\in V_f - S} \mathrm{Gal}(\mathbb{F}/\mathbb{Q}(\Pi'_v)) \\
&\underset{\text{Lem. 8.4}}{=} \bigcap_{v\in V_f - S} \{\sigma \in \mathrm{Gal}(\mathbb{F}/\mathbb{Q}) | {}^{\tilde{\sigma}}\Pi'_v \cong \Pi'_v \text{ for all lifts } \tilde{\sigma} \in \mathrm{Aut}(\mathbb{C})\} \\
&= \bigcap_{v\in V_f} \{\sigma \in \mathrm{Gal}(\mathbb{F}/\mathbb{Q}) | {}^{\tilde{\sigma}}\Pi'_v \cong \Pi'_v \text{ for all lifts } \tilde{\sigma} \in \mathrm{Aut}(\mathbb{C})\},
\end{aligned}$$

where the last line uses Thm. 7.27 and Strong Multiplicity One for discrete series representations of $G'(\mathbb{A})$. Therefore, we get even more that

$$\mathrm{Gal}(\mathbb{F}/\mathbb{E}) = \{\sigma \in \mathrm{Gal}(\mathbb{F}/\mathbb{Q}) | {}^{\tilde{\sigma}}\Pi'_f \cong \Pi'_f \text{ for all lifts } \tilde{\sigma} \in \mathrm{Aut}(\mathbb{C})\}$$

which is obviously the image of $\mathfrak{S}(\Pi'_f)$ under $r_{\mathbb{F}}$. ☐

We may now give the

Proof of Prop. 8.2: Let Π' be regular algebraic and assume that $JL(\Pi')$ is cuspidal. Let $\mathbb{E}$ be the compositum of the fields $\mathbb{Q}(\Pi'_v)$, $v \in V_f - S$. By Thm. 8.1, $\mathbb{Q}(\Pi'_f)$ is a number field. Let $\mathbb{F}$ be the Galois closure of $\mathbb{Q}(\Pi'_f)$. This is a finite Galois extension of $\mathbb{Q}$. We get

$$\begin{aligned}
\mathbb{Q}(\Pi'_f) &= \{z \in \mathbb{C} | \quad \sigma(z) = z \quad \forall \sigma \in \mathfrak{S}(\Pi'_f)\} \\
&= \{z \in \mathbb{F} | \quad \sigma(z) = z \quad \forall \sigma \in r_{\mathbb{F}}(\mathfrak{S}(\Pi'_f))\} \\
&\underset{\text{Lem. 8.5}}{=} \{z \in \mathbb{F} | \quad \sigma(z) = z \quad \forall \sigma \in \mathrm{Gal}(\mathbb{F}/\mathbb{E})\} \\
&= \mathbb{E}.
\end{aligned}$$

☐

Theorem 8.6. *Let $\Pi' \in \mathscr{D}(G')$ be regular algebraic and assume that $JL(\Pi')$ is cuspidal. Let E_μ be a highest weight representation of G'_∞ with respect to which Π'_∞ is cohomological. Then Π'_f is defined over the composite field $\mathbb{Q}(\Pi') = \mathbb{Q}(\mu)\mathbb{Q}(\Pi'_f)$. In particular, Π'_f is defined over a number field.*

Proof. By assumption Π'_∞ is cohomological with respect to the highest weight representation E_μ. In particular, by [7], I, Thm. 5.3.(ii), the central character $\omega_{\Pi'_\infty}$ of Π'_∞ and the central character of $E^\vee_\mu$ coincide. Therefore, $\omega_{\Pi'_\infty}$ is defined over the number field $\mathbb{Q}(\mu^\vee)$, cf. Lem. 7.1, and we may henceforth assume without loss of generality that the central character of Π'_∞ and of E_μ is trivial on $\mathbb{R}_+$.

To proceed, let b be the minimal degree in which the $(\mathfrak{g}'_\infty, K'^\circ_\infty)$-cohomology of $\Pi'_\infty \otimes E_\mu$ does not vanish and let r be the number of split real places. The group $K'_\infty/K'^\circ_\infty \cong (\mathbb{Z}/2\mathbb{Z})^r$ acts on the cohomology $H^b(\mathfrak{g}'_\infty, K'^\circ_\infty, \Pi'_\infty \otimes E_\mu)$. As in [29], 3.3, or [25], 3.1.2, we may pick an $\epsilon \in (K'_\infty/K'^\circ_\infty)^*$ such that its isotypic component $H^b(\mathfrak{g}'_\infty, K'^\circ_\infty, \Pi'_\infty \otimes E_\mu)[\epsilon]$ becomes one-dimensional. As in the proof of Prop. 7.20, we may view Π'_f as an irreducible $G'(\mathbb{A}_f)$-submodule of $H^b_!(S_{G'}, \mathcal{E}_\mu)[\epsilon]$ - the ϵ-isotypic component of interior cohomology in degree b. Let $S \subset V_f$ be a finite set of places, such that Π'_f is unramified outside S and put $K'_S := \prod_{v\notin S} GL_n(\mathcal{O}_v)$ and $G'(\mathbb{A}_S) := \prod_{v\notin S} GL_n(F_v)$. For any subfield $\mathbb{F} \subset \mathbb{C}$ let

$$\mathcal{H}(K'_S, \mathbb{F}) := C^\infty_c\left(G'(\mathbb{A}_S) /\!\!/ K'_S, \mathbb{F}\right)$$

be the K'_S-leveled Hecke algebra of $\mathbb{F}$-valued functions. Hence, $H^b_!(S_{G'},\mathcal{E}_\mu)[\epsilon]^{K'_S}$ is a $\mathcal{H}(K'_S,\mathbb{C})$-module and by what we just said, together with Strong Multiplicity One and Multiplicity One for discrete series representations of $G'(\mathbb{A})$, $\Pi_f'^{K'_S}$ appears precisely once as a submodule in $H^b_!(S_{G'},\mathcal{E}_\mu)[\epsilon]^{K'_S}$. In Lem. 7.3, we have established a $\mathbb{Q}(\mu)$-structure $H^b_!(S_{G'},\mathcal{E}_{\mu,\mathbb{Q}(\mu)})$ on interior cohomology. Letting $\mathbb{Q}(\Pi')=\mathbb{Q}(\mu)\mathbb{Q}(\Pi'_f)$ be the compositum of the rationality fields of E_μ and Π'_f, we obtain a $\mathbb{Q}(\Pi')$-structure $H^b_!(S_{G'},\mathcal{E}_{\mu,\mathbb{Q}(\Pi')})$ and hence also a $\mathbb{Q}(\Pi')$-structure $H^b_!(S_{G'},\mathcal{E}_{\mu,\mathbb{Q}(\Pi')})[\epsilon]$. We now claim that the $\mathcal{H}(K'_S,\mathbb{Q}(\Pi'))$-module $H^b_!(S_{G'},\mathcal{E}_{\mu,\mathbb{Q}(\Pi')})[\epsilon]^{K'_S}$ splits as a direct sum

$$H^b_!(S_{G'},\mathcal{E}_{\mu,\mathbb{Q}(\Pi')})[\epsilon]^{K'_S}\cong W_{\Pi'_f}\oplus W',$$

where $W_{\Pi'_f}\otimes_{\mathbb{Q}(\Pi')}\mathbb{C}\cong\Pi_f'^{K'_S}$ as a $\mathcal{H}(K'_S,\mathbb{C})$-module. This may be seen as follows: The Hecke algebra $\mathcal{H}(K'_S,\mathbb{C})$ decomposes as a restricted tensor product

$$\mathcal{H}(K'_S,\mathbb{C})={\bigotimes_{v\notin S}}'\mathcal{H}(K'_v,\mathbb{C}),$$

$K'_v=GL_n(\mathcal{O}_v)$, of local Hecke algebras $\mathcal{H}(K'_v,\mathbb{C}):=C^\infty_c(GL_n(F_v)/\!\!/K'_v,\mathbb{C})$, each of which is spanned by the Hecke operators $T_{v,j}$, $1\leq j\leq n$, represented by the diagonal matrix $T_{v,j}=\mathrm{diag}(\varpi_v,\dots,\varpi_v,1,\dots,1)$ having exactly j-times ϖ_v on the main diagonal. On a normalized spherical vector of Π'_v, $v\notin S$, $T_{v,j}$ operates by multiplication by the scalar $f_j(\alpha_1^{\Pi'_v},...,\alpha_n^{\Pi'_v})$, cf. Sect. 8.2. Therefore, the functional

$$\lambda_{\Pi'_f,S}:\mathcal{H}(K'_S,\mathbb{C})={\bigotimes_{v\notin S}}'\mathcal{H}(K'_v,\mathbb{C})\to\mathbb{C}$$

determined by the local conditions

$$\lambda_{\Pi'_f,S}(T_{v,j})=f_j(\alpha_1^{\Pi'_v},...,\alpha_n^{\Pi'_v})$$

defines an eigenfunction for the action of $\mathcal{H}(K'_S,\mathbb{C})$ on $H^b_!(S_{G'},\mathcal{E}_\mu)[\epsilon]^{K'_S}$. Recalling from Lem. 8.3 that $\mathbb{Q}(\Pi'_v)=\mathbb{Q}\left(f_1(\alpha_1^{\Pi'_v},...,\alpha_n^{\Pi'_v}),...,f_n(\alpha_1^{\Pi'_v},...,\alpha_n^{\Pi'_v})\right)$ and from Prop. 8.2 that $\mathbb{Q}(\Pi'_f)$ is the compositum of the fields $\mathbb{Q}(\Pi'_v)$, $v\notin S$, we obtain that

$$H^b_!(S_{G'},\mathcal{E}_{\mu,\mathbb{Q}(\Pi')})[\epsilon]^{K'_S}\cong W_{\Pi'_f}\oplus W',$$

where $W_{\Pi'_f}$ is the eigenspace of $\lambda_{\Pi'_f,S}$. But this implies by the one-dimensionality of $H^b(\mathfrak{g}'_\infty,K'^\circ_\infty,\Pi'_\infty\otimes E_\mu)[\epsilon]$ that $W_{\Pi'_f}\otimes_{\mathbb{Q}(\Pi')}\mathbb{C}\cong\Pi_f'^{K'_S}$ as a $\mathcal{H}(K'_S,\mathbb{C})$-module, which shows the above claim.

Now, let $V_{\Pi'_f}$ be the $\mathbb{Q}(\Pi')$-span of the $G'(\mathbb{A}_f)$-orbit of $W_{\Pi'_f}$ inside the $G'(\mathbb{A}_f)$-module $H^b_!(S_{G'},\mathcal{E}_{\mathbb{Q}(\Pi')})[\epsilon]$. In other words, if $\mathcal{H}(G'(\mathbb{A}_f),\mathbb{Q}(\Pi')):=C^\infty_c(G'(\mathbb{A}_f),\mathbb{Q}(\Pi'))$, then

$$V_{\Pi'_f}=\mathcal{H}(G'(\mathbb{A}_f),\mathbb{Q}(\Pi'))\cdot W_{\Pi'_f}.$$

We can finish the proof of the first assertion of theorem, if we show that the natural map

$$V_{\Pi'_f}\otimes_{\mathbb{Q}(\Pi')}\mathbb{C}\to\Pi'_f$$

is an isomorphism. Injectivity follows from the following consideration:

$$\begin{aligned}
\Pi'_f &= \otimes_{v\in S}\Pi'_v\otimes\left(\mathcal{H}(G'(\mathbb{A}_S),\mathbb{C})\cdot\left(\otimes'_{v\notin S}\Pi'_v\right)^{K'_S}\right)\\
&= \mathcal{H}(G'(\mathbb{A}_f),\mathbb{C})\cdot\left(\Pi_f'^{K'_S}\right)\\
&\cong \mathcal{H}(G'(\mathbb{A}_f),\mathbb{C})\cdot\left(W_{\Pi'_f}\otimes_{\mathbb{Q}(\Pi')}\mathbb{C}\right)\\
&\supseteq \left(\mathcal{H}(G'(\mathbb{A}_f),\mathbb{Q}(\Pi'))\cdot W_{\Pi'_f}\right)\otimes_{\mathbb{Q}(\Pi')}\mathbb{C}\\
&= V_{\Pi'_f}\otimes_{\mathbb{Q}(\Pi')}\mathbb{C};
\end{aligned}$$

while surjectivity is a consequence of the irreducibility of Π'_f. Hence, $V_{\Pi'_f}\otimes_{\mathbb{Q}(\Pi')}\mathbb{C}\cong\Pi'_f$.

Recalling Lem. 7.1 and Thm. 8.1 shows that $\mathbb{Q}(\Pi')$ is a number field. Hence, also the second assertion of the theorem follows and the proof is complete. $\square$

Finally, let us investigate the interplay between rationality fields and the global Jacquet-Langlands transfer. Doing so, we get the following Proposition.

Proposition 8.7. *Let* $\Pi' \in \mathscr{D}(G')$ *be regular algebraic and assume that* $JL(\Pi')$ *is cuspidal. Then*

$$\mathbb{Q}(\Pi'_f) = \mathbb{Q}(JL(\Pi')_f).$$

Proof. Let Π' be regular algebraic such that $JL(\Pi')$ is cuspidal. Let S be the finite set of non-archimedean places $v \in V_f$ where Π'_v ramifies. By Cor. 5.5, $\mathbb{Q}(\Pi'_v) = \mathbb{Q}(JL(\Pi')_v)$ for all $v \in V_f - S$. The proposition now follows from Prop. 8.2. $\square$

Remark 8.8. Prop. 8.7 re-proves Thm. 8.1 using the fact that – under the present assumptions – $\mathbb{Q}(JL(\Pi'_f))$ is a number field. (The latter is implicit in Clozel [8], Thm. 3.13.)

8.3. The results of this paper suggest the following

Schólion. *Assuming the cuspidality of* $JL(\Pi')$ *– and not of* Π' *itself – is the appropriate setup to generalize statements on the arithmetic properties of cuspidal automorphic representations from* GL_n/F *to its inner forms* GL'_m/F.

This *schólion* is in accordance with the results of Waldspurger in the case $m = 1$ and $d = 2$, [41]: His space "$A_0(h)$" consists of cohomological cuspidal automorphic representations Π' of $GL'_1(\mathbb{A})$, which are not characters. But for GL_1 over a quaternion division algebra, the latter condition is equivalent to lifting to a cuspidal automorphic representation $JL(\Pi')$ of $GL_4(\mathbb{A})$.

In view of this Schólion we may generalize Clozel's [8, Conjectures 3.7 and 3.8] as:

Conjecture 8.9. *Let* $\Pi' \in \mathscr{D}(G')$ *such that* $JL(\Pi')$ *is cuspidal. Then the following are equivalent:*

(i) Π'_f *is defined over a number field.*
(ii) Π' *is algebraic.*

References

[1] I. A. Badulescu, Global Jacquet-Langlands correspondence, multiplicity one and classification of automorphic representations (with an appendix by N. Grbac), *Invent. Math.* **172** (2008) 383–438

[2] I. A. Badulescu, D. Renard, Unitary Dual of GL(n) at archimedean places and global Jacquet-Langlands correspondence, *Comp. Math.* **145** (2010) 1115–1164

[3] A. Borel, Automorphic Forms on Reductive Groups, in *Automorphic forms and Applications*, IAS/Park City Mathematics Series, vol. 12, eds. P. Sarnak and F. Shahidi, (Utah, 2002) AMS, 2007, pp. 5–40.

[4] A. Borel, Some finiteness properties of adele groups over number fields, *Publ. Math. IHES* **16** (1963) pp. 5–30

[5] A. Borel, H. Jacquet, *Automorphic forms and automorphic representations*, in: Proc. Sympos. Pure Math., Vol. XXXIII, part I, AMS, Providence, R.I., (1979), pp. 189-202

[6] A. Borel, J.-P. Serre, Corners and arithmetic groups, *Comm. Math. Helvet.* **48** (1973) 436–491

[7] A. Borel, N. Wallach, *Continuous cohomology, discrete subgroups and representations of reductive groups*, Ann. of Math. Studies, (Princeton Univ. Press, New Jersey, 2000)

[8] L. Clozel, Motifs et Formes Automorphes: Applications du Principe de Fonctorialité, in *Automorphic forms, Shimura varieties, and L-functions*, Vol. I, Perspect. Math., vol. 10, eds. L. Clozel and J. S. Milne, (Ann Arbor, MI, 1988) Academic Press, Boston, MA, 1990, pp. 77–159.

[9] L. Clozel, On the cohomology of Kottwitz's arithmetic varieties, *Duke Math. J.* **72** (1993) pp. 757–795

[10] P. Deligne, D. Kazhdan, M.-F. Vignéras, *Représentations des algèbres centrales simples p-adiques*, in: Representations of reductive groups over a local field, Travaux en Cours, Hermann, Paris, (1984), pp. 33–117

[11] J. Franke, Harmonic analysis in weighted L_2-spaces, *Ann. Sci. de l'ENS* 4^e série tome 31 **2** (1998) 181–279

[12] J. Franke, J. Schwermer, A decomposition of spaces of automorphic forms, and the Eisenstein cohomology of arithmetic groups, *Math. Ann.*, **311** (1998), pp. 765-790

[13] S. Gelbart, H. Jacquet, Forms on $GL(2)$ from the analytic point of view, in: *Proc. Sympos. Pure Math.*, Vol. XXXIII, part I, AMS, Providence, R.I., (1979), pp. 213–251
[14] H. Grobner, Automorphic Forms, Cohomology and CAP Representations. The Case GL_2 over a definite quaternion algebra, `arXiv:1003.2326v3` preprint (2010)
[15] G. Harder, Eisenstein Cohomology of Arithmetic Groups. The Case GL_2, *Invent. Math.* **89** (1987) pp. 37–118
[16] G. Harder, *General aspects in the theory of modular symbols*, in: *Séminaire de Théorie des Nombres, Paris 1981-82*, ed M.-J. Bertin, Progress in Mathematics **38** (Boston, Basel, Stuttgart 1983), pp. 73-88
[17] Harish-Chandra, Harmonic Analysis on Real Reductive Groups III, *Ann. Math.* **104** (1976) 117–201
[18] M. Harris, R. Taylor *The geometry and cohomology of some simple Shimura varieties*, Annals of Math. Studies **151**, Princeton Univ. Press (2001)
[19] T. W. Hungerford, *Algebra*, Graduate Texts in Mathematics, Springer (2003)
[20] H. Jacquet, R. P. Langlands, *Automorphic forms on $GL(2)$*, Lecture Notes in Mathematics, No. **114**, Springer-Verlag (1970)
[21] A. W. Knapp, D. A. Vogan Jr. *Cohomological induction and unitary representations*, Princeton Univ. Press, (1995)
[22] R. P. Langlands, *On the Classification of Irreducible Representations of Real Algebraic Groups*, Mathematical Surveys and Monographs, AMS, (1988)
[23] R. P. Langlands, *On the functional equations satisfied by Eisenstein series*, Lecture Notes in Mathematics **544** (1976)
[24] J.-S. Li, J. Schwermer, On the Eisenstein Cohomology of Arithmetic Groups, *Duke Math. J.* **123** (2004) pp. 141–169
[25] J. Mahnkopf, Cohomology of arithmetic groups, parabolic subgroups and the special values of automorphic L-Functions on $GL(n)$, *Journal de l'Institut de Mathématiques de Jussieu* **4** (2005) pp. 553–637.
[26] C. Mœglin, J.-L. Waldspurger, Le Spectre Résiduel de $GL(n)$, *Ann. Sci. de l'ENS* 4^e série tome 22 **4** (1989) 605–674
[27] C. Mœglin, J.-L. Waldspurger, *Spectral decomposition and Eisenstein series*, Cambridge Univ. Press (1995)
[28] A. Raghuram, F. Shahidi, Functoriality and special values of L-functions, in: *Eisenstein series and Applications*, eds W. T. Gan, S. Kudla, Y. Tschinkel, Progress in Mathematics **258** (Boston, 2008), pp. 271–294.
[29] A. Raghuram, F. Shahidi, On certain period relations for cusp forms on GL_n, *Int. Math. Res. Not.* (2008) Art. ID rnn 077, 23 pp.
[30] J. G. Rogawski, Representations of GL(n) and division algebras over a p-adic field, *Duke Math. J.* **50** (1983) 161–196
[31] J. Rohlfs, Projective limits of locally symmetric spaces and cohomology, *J. reine angew. Math.* **479** (1996) 149–182
[32] S. A. Salamanca-Riba, On the unitary dual of real reductive Lie groups and the $A_{\mathfrak{q}}(\lambda)$ modules: The strongly regular case, *Duke Math. J.* **96** (1998) 149–159
[33] J. Schwermer, *Cohomology of arithmetic groups, automorphic forms and L-functions*, in: *Cohomology of Arithmetic Groups and Automorphic Forms*, eds. J.-P. Labesse, J. Schwermer, Lecture Notes in Math. **1447** (Springer, 1990) pp. 1–29
[34] J. A. Shalika, The multiplicity one theorem for GL_n, *Ann. Math.* **100** (1974) pp. 171-193
[35] G. Shimura, The special values of the zeta functions associated with Hilbert modular forms, *Duke Math. J.* 45 (1978), no. 3, 637–679.
[36] B. Speh, Unitary representations of $GL(n,\mathbb{R})$ with nontrivial $(\mathfrak{g},K)$-cohomology, *Invent. Math.* **71** (1983) 443-465
[37] J. Tate, *Number Theoretic Background*, in: Proc. Sympos. Pure Math., Vol. XXXIII, part II, AMS, Providence, R.I., (1979), pp. 3–26
[38] D. A. Vogan Jr., Gelfand-Kiriliov Dimension for Harish-Chandra Modules, *Invent. Math.* **48** (1978) 75–98
[39] D. A. Vogan Jr., Unitarizability of certain series of representations, *Ann. of Math.* **120** (1984) pp. 51-90
[40] D. A. Vogan Jr., G. J. Zuckerman, Unitary representations with nonzero cohomology, *Comp. Math.* **53** (1984) 51–90
[41] J.-L. Waldspurger, Quelques propriétés arithmétiques de certaines formes automorphes sur GL(2), *Comp. Math.* **54** (1985) 121–171
[42] N. Wallach, *On the constant term of a square integrable automorphic form*, in: Operator algebras and group representations, Vol. II (Neptun, 1980), Monographs Stud. Math., vol. **18**, Pitman, Boston, MA, London (1984), pp. 227-237

HARALD GROBNER: *Current address*: INSTITUT DE MATHÉMATIQUES DE JUSSIEU, 4, PLACE JUSSIEU, 75252 PARIS CEDEX 05, FRANCE.
E-mail address: `harald.grobner@univie.ac.at`

A. RAGHURAM: DEPARTMENT OF MATHEMATICS, OKLAHOMA STATE UNIVERSITY, 401 MATHEMATICAL SCIENCES, STILLWATER, OK 74078, USA.
E-mail address: `araghur@math.okstate.edu`